

# Noxious WEED CONTROL

FS525N



Mike Moechnig, Extension Weed Specialist  
Leon J. Wrage, Extension Emeritus, Plant Science  
Darrell L. Deneke, Extension IPM Program

## WATCH for NOXIOUS WEED INVADERS

Noxious weeds are non-native plant species that are a concern to South Dakota land owners and managers. They can seriously impact the native plant community by altering or affecting agriculture, recreation, and wildlife.

### 2006 Noxious Weed Estimates Statewide

	<i>Acres*</i>
Canada thistle	1,699,994
Leafy spurge	316,671
Perennial sowthistle	115,105
Russian knapweed	2,372
Hoary cress	10,936
Purple loosestrife	1,742
Saltcedar	6,273

\* Estimates from 2006 State Noxious Weed Annual Report, SD Department of Agriculture.

## LOCAL NOXIOUS WEEDS

Several weeds have been designated locally noxious in one or more counties by the South Dakota Weed and Pest Commission. Statute requirements for control are the same as for statewide noxious weeds.

- absinth wormwood
- bull thistle
- chicory
- common burdock
- common mullein
- common tansy
- dalmatian toadflax
- diffuse knapweed
- field bindweed
- giant knotweed
- houndstongue
- musk thistle
- plumeless thistle
- poison hemlock
- puncturevine
- Scotch thistle
- spotted knapweed
- St. Johnswort
- yellow toadflax

## NOXIOUS WEED OF CONCERN

Two invasive species are listed as “local noxious weeds” by the South Dakota Weed and Pest Control Commission. They are yellow and Dalmatian toadflax.

Introduced to the U.S. as ornamental garden plants—the flowers resemble snapdragon blooms and yellow toadflax is commonly known as butter and eggs—both species spread aggressively. Grass productivity can be reduced significantly in infested areas, and both toadflaxes are considered to be unpalatable to wildlife and cattle.

Prevention is the best control. Both yellow and Dalmatian toadflax have extensive root systems that make them competitive and difficult to control once they get a start in the field. Maintain the health of desirable pasture plants and avoid overgrazing; toadflaxes like disturbed, overgrazed areas. Herbicide options are available; timing of application depends on the species.



**Mature Dalmatian toadflax**



**Mature yellow toadflax**

**SOUTH DAKOTA STATE UNIVERSITY**

College of Agriculture & Biological Sciences • Cooperative Extension Service • U.S. Department of Agriculture

## CONTENTS

---

2,4-D Rate - Product Chart . . . . .	3	Houndstongue . . . . .	20
Glyphosate Product Chart . . . . .	3	Toadflax (Dalmatian and yellow) . . . . .	21
Dicamba Product Chart . . . . .	3	Puncturevine . . . . .	22
Abbreviations . . . . .	4	Poison Hemlock . . . . .	22
Leafy Spurge, Hoary Cress, Field Bindweed . . . . .	4	Saltcedar . . . . .	23
Canada Thistle, Perennial Sow Thistle . . . . .	6	Shelterbelts . . . . .	24
Biennial Thistles (musk, plumeless, bull, Scotch) . . . . .	10	Cultivated Cropland . . . . .	25
Absinth Wormwood (wormwood sage) . . . . .	13	Aquatic . . . . .	27
Knapweed Species (Russian, spotted, diffuse) . . . . .	14	Noncrop and Special Situations . . . . .	28
Mullein . . . . .	16	2,4-D Label Restrictions & Noncrop Labeling . . . . .	30
Purple Loosestrife . . . . .	17	Summary Site and Use Restrictions . . . . .	31
Tansy . . . . .	18	Chemical, Physical, and Safety Characteristics of Herbicides . . . . .	33
St. Johnswort . . . . .	19	Biocontrol of Noxious Weed . . . . .	34

---

**This publication and others can be accessed electronically from the SDSU  
College of Agriculture & Biological Sciences publications page, which is at  
<http://agbiopubs.sdstate.edu/articles/FS525N.pdf>**



South Dakota State University, South Dakota counties, and U.S. Department of Agriculture cooperating. South Dakota State University is an Affirmative Action/Equal Opportunity Employer and offers all benefits, services, education, and employment opportunities without regard for race, color, creed, religion, national origin, ancestry, citizenship, age, gender, sexual orientation, disability, or Vietnam Era veteran status.

6,800 copies printed by CES at a cost of ? each. FS525N. February 2007

## HERBICIDES for NOXIOUS WEEDS

**Noxious Weed Recommendations:** Herbicides for pasture, range, and noncrop areas, including roadside and other rights-of-way that may be harvested for hay or grazed, are given a priority.

**Noncrop Areas.** Noncropland is defined for herbicide purposes as areas not used to produce food or feed crops during the time herbicides residue remains in the soil. Noncrop areas include parking lots, utility storage areas and some rights-of-way. Pasture, range, and hayland are cropland.

**Herbicides.** Herbicides are listed by tradename except where the active ingredient is available in several products. The common name (in parentheses) follows the first listing of the tradename. Product labels for the same active ingredient vary. Herbicides included are those considered for most situations and those generally available. Rights-of-way are frequently grazed or used for hay, therefore products that allow grazing or harvesting hay are given a priority.

Rates are based on research data available. Treatments include those that (1) provide a high level of eradication with one application for small areas; (2) have lower annual cost but give a high degree of eradication in a four or five year program or (3) have low cost but require repeated application for several years.

*Tradenames for herbicides are used in this publication to aid reader recognition. The common name is also listed and is used for herbicides that are available in many labeled products. Examples of other product names are listed where possible based on information available. As patents expire and marketing agreements are formed, additional products may be marketed. Be sure crop use and application directions are followed for the product being used.*

Rates for 2,4-D are stated as acid equivalent (ai) per acre. The amount of product for several rates is listed for each formulation.

<u>Lb/A Required</u>	<b>2,4-D RATE Product Per Acre</b>		<u>FORMULATION</u>	
	<u>3.8L*</u>	<u>5.7L*</u>	<u>80% WSP</u>	<u>90% WSP</u>
0.5	1 pt	.66 pt	.66 lb	.6 lb
1.0	2 pt	1.33 pt	1.25 lb	1.1 lb
1.5	3 pt	2 pt	1.9 lb	1.7 lb
2.0	4 pt	2.66 pt	2.5 lb	2.2 lb

\*2,4-D showing 3.8 lb/gal is the same as 4 lb/gal; and 5.7 lb/gal is the same as 6 lb/gal acid equivalent.

### GLYPHOSATE PRODUCTS Equivalent Rates

Glyphosate is formulated in different salts and different concentrations. Formulation concentration is listed as lb of glyphosate acid/gal (ae); some labels also list lb glyphosate acid plus salt/gal (active ingredient, ai).

<u>Formulation</u>		<u>Amount of Product for lb ae</u>			
		<u>.38 ae</u>	<u>.75 ae</u>	<u>1.5 ae</u>	<u>3 ae</u>
3 lb ae (4 lb ai)	L	1 pt	2 pt	2 qt	4 qt
3.7 lb ae (5 lb ai)	L	.8 pt	1.6 pt	1.6 qt	3.2 qt
4 lb ae (5.4 lb ai)	L	.75 pt	1.5 pt	1.5 qt	3 qt
4.17 lb ae	L	.75 pt	1.5 pt	1.5 qt	3 qt
4.5 lb ae (5.5 lb ai)	L	.7 pt	1.4 pt	1.4 qt	2.8 qt
65% ae (72% ai)	DS	.6 lb	1.1 lb	2.3 lb	4.6 lb

### DICAMBA PRODUCTS Formulations

<u>Product</u>	<u>lb a.e./gal</u>	<u>Formulation</u>
Banvel	4	dimethylamine salt
Clarity	4	diglycolamine salt
Sterling	4	dimethylamine salt
Vanquish	4	diglycolamine salt

**ABBREVIATIONS.** Several abbreviations are used in this publication.

pt = pint  
qt = quart  
gal = gallon  
lb = pound  
oz = ounce  
t = teaspoon  
T = Tablespoon

ae = acid equivalent  
ai = active ingredient  
L = liquid  
gpa = gallons per acre  
psi = lb per square inch  
DF = dry flowable  
DS = dry soluble  
WDG = water soluble powder or crystals  
DG = dispersible granule  
NIS = non-ionic surfactant  
XP = extruded paste

## LEAFY SPURGE, HOARY CRESS, and FIELD BINDWEED

---

### **Tordon 22K (picloram) Leafy spurge and field bindweed**

**1-2 qt Tordon 2L (1 lb ae)**

**(\$46.25)**

**Spring or Fall.** Primarily for small patches. Use 2 qt/A Tordon as a spot treatment. Rates above 1 qt/A cannot exceed 50% of an acre. Rates above 1 pt/A can be used in fallow cropland if the treated areas are less than 10% of the field. Tordon is an effective choice for leafy spurge. Minimum carrier is 20 gpa. The 2 qt rate will reduce the stand; however, additional Tordon or other follow-up is required. Some regrowth may be noted after application during dry seasons. Make spring treatments before seeds form. Make fall application before soil freeze-up.

Tordon has foliar activity and extended soil residual. It controls topgrowth and translocates into roots. Rainfall is required to move the herbicide into the root zone. Trees, legumes, and broadleaf plants are very sensitive to drift and soil residues.

Tordon is registered for use in grass pasture and range, fallow cropland and noncrop areas. Bromegrass, buffalograss, and wheatgrass may be injured; bluegrass is tolerant. Tordon is formulated as a liquid containing 2 lb/gal acid equivalent.

**RESTRICTIONS:** Do not use in trees. Avoid drift to trees or sensitive broadleaf crops. For rates above 1 qt/A, do not harvest for hay within 2 weeks after treatment. Do not graze dairy animals for 2 weeks after treatment. Remove animals 3 days before slaughter if grazing within 2 weeks after spraying. Do not apply into water or wetlands or on inner banks of irrigation or drainage ditches. Risk of leaching is greatest where soils have rapid permeability (such as loamy to sand) and where the underlying aquifer is near the surface. **Restricted Use Pesticide.**

---

### **Tordon 22K + 2,4-D Ester (picloram + 2,4-D) Leafy spurge and field bindweed**

**1.5 pt Tordon 2L + 1 lb ae 2,4-D ester (.38 + 1 lb ae)**

**(\$21.25)**

**Spring.** Tank-mix. Intended as a 4 to 5 year program. For leafy spurge, apply in late bud stage. Use Tordon at 1.5 pt plus 2,4-D ester at 1 lb ae/A. Lower rates may not provide sufficient residual control into the fall. Amine formulation of 2,4-D may be used if site limitations preclude ester formulations. Minimum carrier is 10 gpa for ground or 2 gpa for air. Intended as one application per year; some regrowth may be noted in wet seasons. Treatment has provided 75 to 80% leafy spurge stand reduction after 4 years.

**RESTRICTIONS:** Do not use in trees. Avoid drift to sensitive broadleaf crops. Do not graze lactating dairy for 7 days after application. Labels for 2,4-D allow harvesting hay 30 days after application and require a 3 day removal period before slaughter. Refer to surface or groundwater precautions as for Tordon alone. **Restricted Use Pesticide.**

---

### **Plateau (imazapic) Leafy spurge, hoary cress, and field bindweed**

**8-12 oz Plateau 2L (.12-.18 lb ai)**

**(\$18.20-27.25)**

**Fall.** Plateau is an imidazolinone herbicide used to control leafy spurge in pasture, range, noncrop areas, and CRP plantings. Plateau is being marketed only through public agencies and organizations. However, it is labeled and may be applied to private as well as public land.

Results have been promising in SDSU tests; 70 to 90% control has been reported the year after application. Fall treatment to active growth has been more effective than spring applications. Plateau is available as a 2 lb/gal liquid.

## PLATEAU (Continued . . . )

Suggested rates are 8 to 12 oz/A of 2L. Use the higher rate for dense infestations that have been established for longer periods of time. The lower rate has been used in most tests. Field plot data suggest follow-up control programs are required to prevent reinfestation. Add MSO at 2 pt plus 2 pt/A 28% N. Minimum carrier is 10 gpa for conventional ground equipment or 2 gpa for low volume equipment.

Grass tolerance is an important issue. Plateau is labeled as a seedhead growth suppressant for cool-season grasses. It also is labeled for range restoration to release native warm-season species such as bluestem, needlegrass, and Indiangrass using the same rates. It may also be used at 4 oz/A on certain species in new CRP seedings.

Several tree and shrub species listed on the label are known to have acceptable tolerance when applied under the canopy and/or to the foliage. Tolerance is based upon trees with a minimum of 2 inch DBH (diameter at breast height). Some species may exhibit tip chlorosis and minor necrosis. Foliar contact on some species may increase injury, defoliation, and terminal death.

**RESTRICTIONS:** Treated areas may be grazed. Do not harvest hay for 7 days after treatment.

---

## 2,4-D Ester *Leafy spurge and field bindweed*

**1.5 lb ae 2,4-D ester 4L (1.5 lb ae) (\$5.85)**

**Spring** and **Fall.** Spring and fall application required each year. Selective, foliage applied, translocated herbicide. Uses for 2,4-D include grass pasture, range, and noncrop areas. Apply 2,4-D ester at 1.5 lb ae/A. Suggested carrier is 10 to 40 gpa. This treatment has been popular for large infestations; however, several years are required to achieve significant stand reduction.

Apply in spring at late bud stage when bracts begin to yellow. Retreat in September or early October when new fall growth is 4 to 6 inches. Results can be variable. One treatment per year prevents seed production. Stands have been reduced 50% in 3 years. Complete eradication is difficult even after 10 years. Surfactant or fuel additives increase leaf burn but seldom increase stand reduction.

**Spring** or **Fall.** Single application each year. Rates to 2 lb/A may be used in pasture and range; higher rates are allowed in noncrop and fallow. Control is less than for two applications of 1.5 lb/A each. Best for inaccessible areas where the labor cost for a second application is prohibitive.

**RESTRICTIONS:** Avoid drift to trees and sensitive crops. Not suggested for use in trees. Do not graze lactating dairy animals for 7 days after application. Labels for 2,4-D allow harvesting hay 30 days after application and require a 3 day removal period before slaughter. Note other label precautions. Labels for 2,4-D products vary.

---

## Dicamba Products (*dicamba*) *Field bindweed*

**2 qt dicamba 4L (2 lb ae) (\$32.00)**

**Spring** or **Fall.** Minimum carrier volume is 3 gpa for ground applications or 2 gpa for air. Apply in fall prior to a killing frost. Registered for use in pasture/rangeland and non-crop areas. May stunt brome grass but bluegrass and several other grasses are tolerant. Trees, legumes, and broadleaf plants are sensitive to drift and soil residues.

**RESTRICTIONS:** Do not graze lactating dairy for 40 days or harvest hay for 70 days after application of more than 1 qt/A. Slaughter animals must be removed for 30 days after last application. Avoid drift to non-target plants.

---

## Overdrive (*diflufenzopyr + dicamba*) *Field bindweed*

**4-8 oz Overdrive 70DF (0.05-1 + 0.125-0.25) (\$10.45-20.90)**

**Spring** or **Fall:** Use higher recommended rates for perennial weed species. A maximum of 10 oz/A per year may be applied in non-crop sites or 8 oz/A per year in pasture/range or hay. Use NIS at 1 qt per 100 gallons or MSO at 1.5-2 pt/A. Minimum carrier is 3 gpa. Rainfast 4 hours after application. May be tank-mixed with several other registered herbicides to control additional weed species.

**RESTRICTIONS:** Supplemental label for aerial application in CRP. Do not apply to newly seeded grass. No restrictions for grazing or haying.

---

---

**Telar (chlorsulfuron) Hoary cress****.5-1.3 oz Telar 75XP (.023-.061 lb ai)****(\$10.25-26.60)**

**Spring.** Apply .5 to 1 oz/A of Telar 75DF to hoary cress at bud to bloom stage in spring or in fall at rosette stage. Telar is labeled for use on noncrop, rights-of-way, pasture, range, and CRP. Several wheatgrasses, bluegrass, and smooth brome grass are more tolerant to rates over .5 oz/A; bluestem, buffalograss, green needlegrass, Indiangrass, and switchgrass are tolerant to rates up to .5 oz/A. Minimum carrier is 10 gpa. Add NIS at 1 qt/100 gal.

**RESTRICTIONS:** There are no grazing restrictions for rates less than 1.3 oz/A. Labeling does not include hay uses. Do not apply to water, such as lakes, streams, or areas where runoff flows into such areas.

---

**Escort (metsulfuron) Hoary cress****1-2 oz Escort 60XP (.038-.075 lb ai)****(\$19.65-39.35)**

**Spring.** For hoary cress control in grassland. Apply to actively growing weeds at rosette stage. Escort is for western range, rights-of-way, and noncrop areas. Minimum of 10 gpa carrier is suggested. Add NIS at 1 qt/100 gal.

**RESTRICTIONS:** Legumes will be injured. No haying or grazing restrictions for rates less than 1.66 oz; delay haying for 3 days for higher rates. Do not apply to water, lakes, streams, or areas that may run off into such areas.

---

---

**CANADA THISTLE and PERENNIAL SOW THISTLE**

---

---

**Tordon 22K (picloram)****1-2 qt Tordon 2L (.5-1 lb ae)****(\$23.15-46.25)**

**Spring or Fall.** Primarily for small infestations. Apply 2 qt/A Tordon as a spot treatment. Use a minimum of 20 gpa carrier. Rates above 1 qt/A cannot exceed 50% of an acre. Stands have been reduced 85 to 90% in SDSU tests. Make spring treatments before seed forms. Make fall applications before soil freeze-up. Use 2,4-D or other herbicides to control stragglers or seedlings in succeeding years.

The 1 qt/A rate has been promising in recent Canada thistle tests. Use the high rate for maximum stand reduction, especially for fall treating areas such as fence lines. Control one year later is greater than for lower Tordon rates in tank-mixes.

Tordon has foliar activity and extended soil residual. It is absorbed into the leaves and controls topgrowth. It also translocates into roots. Rainfall moves the herbicide into the root zone. Trees, legumes, and broadleaf plants are very sensitive to drift and soil residual.

Tordon is registered for grass pasture and range, fallow cropland, and noncrop areas. At high rates, brome grass, buffalograss, and wheatgrass may be severely injured; bluegrass is tolerant. Tordon is formulated as a liquid containing 2 lb/gal acid equivalent.

**RESTRICTIONS:** Do not use in trees. Avoid drift to trees or sensitive broadleaf crops. For rates above 1 qt/A, do not harvest for hay within 2 weeks of treatment. Do not graze dairy animals for 2 weeks after treatment. Remove animals 3 days before slaughter if grazing within 2 weeks after spraying. Do not apply into water or wetlands or on inner banks of irrigation or drainage ditches. Risk of leaching is greatest where soils have rapid permeability (such as loamy to sand) and where the underlying aquifer is near the surface. **Restricted Use Pesticide.**

---

---

## Tordon 22K + 2,4-D (*picloram* + 2,4-D)

**1 pt Tordon 2L + 1 lb ae 2,4-D (.25+1 lb ae) (\$15.20)**  
**4 pt Grazon 2.54L (.27+1 lb ae)**

**Spring.** Intended as a 4- to 5-year program with sufficient seasonal suppression so only one application per season is required. Apply at bud stage before flowers open. Tank-mix 1 pt Tordon plus 1 lb ae 2,4-D or use 4 pt Grazon premix containing .54 lb picloram (Tordon) plus 2 lb 2,4-D amine per gal. Amines cause less leaf burn and are preferred if growth is lush. Minimum carrier is 10 gpa for ground 2 gpa for air. This combination is promising to improve stand reduction over a period of several years. Some fall regrowth may be noted in wet seasons. Lower rates do not provide sufficient residual control into the fall.

The tank-mix is registered for noncrop, grass pasture, and range. Grazon may be used only in grass pasture and range. Grass is usually tolerant to these Tordon rates. Some stunting, especially if applied at boot stage, may be noted.

**RESTRICTIONS:** Do not use in trees. Avoid drift to sensitive broadleaf crops. Refer to surface or groundwater precautions as for Tordon alone. Do not graze lactating dairy for 7 days after application. Labels for 2,4-D allow harvesting hay 30 days after application and require a 3 day removal period before slaughter. **Restricted Use Pesticide.**

---

## Milestone (*aminopyralid*)

**3-7 oz Milestone 2L (0.05-0.11 lb ai) (\$7.75-18.10)**

**Spring or Fall.** Will control Canada thistle (5-7 oz/A) and perennial sowthistle (3-5 oz/A). Make applications in early summer before the bud stage or early flowering or in fall prior to a killing frost. Use higher rates for advanced weed growth stages, dense stands, or under adverse growing conditions, such as drought. Use a non-ionic surfactant (NIS) at 0.25-0.5% (1-2 qt per 100 gallons spray solution) under adverse growing conditions or advanced weed growth stages. Avoid mowing for 7 days after application to allow for herbicide translocation in the weeds. SDSU studies have demonstrated excellent control with consistent results.

Milestone may be applied to non-irrigation ditch banks and seasonally dry wetlands, but may not be applied over water or to areas where surface water is present. Applications may be made on warm and cool season grasses.

**RESTRICTIONS:** Do not apply more than 7 fl oz (0.11 lb ae) per acre per year for broadcast applications or 14 fl oz (0.22 lb ae) for spot treatments (less than ½ acre areas). There are no grazing restrictions. Aminopyralid residues may be transferred in manure from livestock that have consumed treated forage within the past 3 days. After grass seeding, wait until perennial grasses are well established with a secondary root system before applying Milestone. Some grasses, such as smooth brome, may be suppressed under adverse growing conditions.

---

## ForeFront (*aminopyralid* + 2,4-D)

**2-2.6 pt ForeFront 3L (0.08-0.11 lb ae aminopyralid + 0.67-0.87 lb ae 2,4-D) (\$14.45-18.75)**

**Spring or Fall:** Apply prior to flowering in early summer or to regrowth in the fall. Minimum recommended carrier volume is at least 10 gpa for ground application or 2 gpa for aerial application. Use a NIS as recommended for Milestone (0.25-0.5% v/v).

**RESTRICTIONS:** Do not harvest forage for hay within 14 days after application. Aminopyralid residues may be transferred in manure from livestock that have consumed treated forage within the past 3 days. Do not apply more than 2.6 pt/A in a growing season.

---

## Stinger or Transline (*clopyralid*)

**.66-1.33 pt Stinger 3L or Transline 3L (.25-.5 lb ae) (\$39.90-80.40)**

**Spring.** For Canada thistle. Perennial sow thistle suppression. Apply from rosette to bud stage when plants are actively growing. Stinger is labeled for use in grass pasture, rangeland, fallow, fencerows, and other noncrop areas. Transline is labeled for noncrop areas, rangeland, and rights-of-way. Transline is marketed only to government and public agencies. Use 1 to 1.33 pt/A to achieve maximum stand reduction. Data suggest 90 to 95% reduction can be expected. Minimum carrier is 5 gpa; use at least 10 gpa for most ground applications.

Stinger or Transline has potential in sites where grass cannot be damaged or where trees limit use of herbicides with harmful soil residual.

**RESTRICTIONS:** Do not contaminate irrigation ditches. No grazing or haying restrictions.

---

## **Curtail (*clopyralid + 2,4-D*)**

**2 qt Curtail 2.38L (.19 + 1 lb ae)**

**(\$18.60)**

**Spring.** Premix. For Canada thistle and perennial sowthistle suppression. Apply before bud stage when plants are actively growing. Reduced rate of 1 qt/A is frequently used for seasonal suppression. The 2 qt/A rate has provided excellent control and 60 to 70% stand reduction. Minimum carrier is 10 gpa for ground and 5 gpa for air for most situations. Curtail is a premix containing .38 lb clopyralid (Stinger) plus 2 lb 2,4-D amine per gallon.

Curtail is labeled for use in noncrop areas, rangeland, grass pasture, and CRP grass seedings and fence lines.

**RESTRICTIONS:** Do not graze lactating dairy cattle in treated areas for 14 days after application. Remove meat animals 7 days before slaughter if grazing within 2 weeks after application. Do not harvest hay within 30 days after application.

---

## **Redeem (*triclopyr + clopyralid*)**

**4 pt Redeem 3L (1.12 + .38 lb ae)**

**(\$47.55)**

**Spring.** Premix. Intended for Canada thistle and certain broadleaf weeds not controlled with 2,4-D. Apply 4 pt/A from early growth to bud stage. Weeds must be actively growing. Reduced rates provide seasonal suppression. Minimum carrier is 10 gpa for ground or 3 gpa for air. Redeem is a premix containing 2.25 lb triclopyr (Garlon) plus .75 lb clopyralid (Stinger) per gallon. Redeem is labeled for use in rangeland, grass pasture, roadsides, CRP grass seedings, fence lines, and other noncrop areas.

**RESTRICTIONS:** Do not graze or harvest forage for dairy cattle for 14 days after application. Do not harvest hay until the next growing season for dairy or harvest hay for other livestock for 7 days after treatment. Remove meat animals at least 3 days before slaughter. Do not treat banks or bottom of irrigation ditches.

---

## **Overdrive (*diflufenzopyr + dicamba*)**

**4-8 oz Overdrive 70DF (.05-.1 + .125-.25 lb ae)**

**(\$10.45-20.90)**

**Spring or Fall.** Rate is 4 to 8 ounces per acre based on weed species and maturity. Use higher rates for perennial weeds like Canada thistle and sowthistle. A maximum of 10 oz/A can be applied per season in noncropland sites and a maximum of 8 ounces per acre in pasture, hay, and rangeland. Use 1 qt NIS per 100 gal or MSO at the rate of 1.5 to 2 pt/A. Do not use less than 3 gallons of spray volume per acre. Rainfast 4 hours after application. Overdrive may be tank-mixed with several labeled tank-mix partners to improve weed control spectrum.

**RESTRICTIONS:** Supplemental label for aerial application in CRP. Do not plant crops for 30 days after last application. Pasture or rangeland grass treated with Overdrive can be grazed or harvested for livestock feed immediately after application. Do not apply to newly seeded grasses or small grains.

---

## **Dicamba Products (*dicamba*)**

**2 qt dicamba 4L (2 lb ae)**

**(\$32.00)**

**Spring or Fall:** Spot treatment rate, primarily for small infestations. Use at least 3 gpa carrier for ground or 2 gpa for air. Make spring application at early bud stage. Apply in fall before a killing frost.

Stand reduction may range from 50-70% the year following application. Lower rates provide less permanent reduction. Control is usually greater in low rainfall areas. An application of 2,4-D or other herbicide is required to control seedlings and weakened plants in succeeding years. Refer to Dicamba Products table. Clarity and Vanquish contain a low vapor salt of dicamba.

Dicamba is a selective, translocated herbicide. It has foliar activity. Favorable growing conditions improve results. High rates have soil residual extending through one season. Dicamba products are registered for use in pasture, range, and noncrop areas. Bromegrass may be severely stunted; bluegrass and several other grasses are tolerant. Trees, legumes, and broadleaved plants are sensitive to drift and soil residues.

**RESTRICTIONS:** Do not graze lactating dairy for 40 days or harvest hay for 70 days after application of more than 1 qt/A. Slaughter animals must be removed for 30 days after last application. Avoid drift to non-target plants. Do not contaminate water.



---

## Dicamba Products + 2,4-D (*dicamba + 2,4-D*)

2 pt dicamba 4L + 1 lb ae 2,4-D (1 + 1 lb ae)

(\$18.90-19.80)

**Spring.** Tank-mix. Intended as a 4- to 5- year program with sufficient seasonal suppression so only one application per year is required. Apply at bud stage. Mix dicamba 4L at 2 pt plus 2,4-D at 1 lb ae/A. Amines cause less leaf burn and are preferred if growth is lush. Apply in 10 to 20 gpa carrier. This combination is a promising program to improve stand reduction over a period of several years. It also is a good choice for inaccessible areas where the labor cost of a second application is prohibitive. Lower dicamba rates may not provide sufficient residual control into the fall, especially in wet seasons. Refer to Dicamba Products table.

Dicamba plus 2,4-D is labeled for use in grass pasture, range, and noncrop areas. Grass is usually tolerant to these rates; some stunting may be noted, especially if applied at boot stage.

**RESTRICTIONS:** Do not use in trees. Avoid drift to sensitive broadleaved plants. Labels for 2,4-D do not allow grazing lactating dairy for 7 days after application. Labels for 2,4-D allow harvesting hay 30 days after application and require a 3 day removal period before slaughter. Dicamba labeling for 1 to 2 pt/A restricts grazing lactating dairy for 21 days or haying for 51 days after application. Note other label restrictions for higher rates.

---

## Telar (*chlorsulfuron*)

1-1.3 oz Telar 75XP (.046-.061 lb ai)

(\$20.50-26.70)

**Spring or Fall.** For Canada thistle. Apply 1.3 oz Telar XP at bud to bloom or in fall at rosette stage. Spring application preferred. Minimum carrier is 10 gpa. Add NIS at 1 qt/100 gal. May be tank-mixed with 2,4-D, dicamba, or glyphosate products.

**RESTRICTIONS:** There are no grazing restrictions for rates less than 1.3 oz/A. Labeling does not include hay uses. Do not apply to water, such as lakes, streams, or areas where runoff flows into such areas.

---

## Cimarron Max (*metsulfuron + dicamba + 2,4-D*)

20-5 Acres/.5 oz Part A + 2.5 gal Part B (.01-.038 + .12-.5 + .38-1.5 lb ae)

(\$6.70-26.80)

Cimarron Max herbicide is a twin-pak combination. Part A contains metsulfuron (Ally). Part B contains 1 lb dicamba (Banvel) plus 2.87 lb ae 2,4-D amine per gallon. The use ratio is .5 oz Part A to 2.5 gal of Part B to treat 5 to 20 acres. Refer to following rate table for acres treated.

---

Cimarron Max Rate	Part A Rate (oz/A)	Part B Rate (pt/A)	Acres Treated with .5 oz Part A + 2.5 gal Part B
Rate I	0.25	1	20A
Rate II	0.5	2	10A
Rate III	1	4	5A

---

Use Rate II for suppression of Canada thistle and Rate III for suppression of perennial sowthistle. The Rate II or 10 acre rate provides the equivalent of .5 oz Ally 60DF + .5 pt Banvel 4L + 1.5 pt 2,4-D 3.8L per acre. Rates are based on weed species and weeds less than 4 inches tall. Do not apply more than the equivalent of 1.66 oz per acre of Cimarron Max Part A per year. Add NIS at 2 to 4 pt/100 gal. May be applied by ground or air. No restrictions or waiting period between treatment and grazing for non-lactating animals. Remove meat animals 30 days prior to slaughter. Do not graze lactating dairy within 7 days of treatment. Do not harvest hay for 37 days after treatment.

---

---

## 2,4-D

### 2-3 lb ae 2,4-D Amine (2-3 lb ae)

**(\$5.80-8.70)**

**Late Fall:** Apply a high rate in late fall before leaves are damaged by frost. Rates to 2 lb ae/A may be used in pasture and range; higher rates are allowed in noncrop and fallow. Data indicate 50 to 60% stand reduction if there is considerable new growth and if weeds have been weakened by previous control practices. Light frost before application does not reduce control; temperatures of 60° F. after application improve results.

### 1.5 lb ae 2,4-D Amine or Ester (1.5 lb ae)

**Spring or Fall:** Requires a spring and fall application each year. Make spring applications at bud stage. Retreat in September or early October after new fall growth reaches 6 inches. Suggested carrier is 10 to 20 gpa. Results can be variable. One spraying prevents seed production. Two applications provided 10 to 25% stand reduction the first year in SDSU tests. Reductions of 70 to 80% may be achieved after 3 years.

Uses for 2,4-D include grass pasture, range, and noncrop areas. This is a popular program for large infestations in pasture and roadsides. However, several years are required to reach high levels of eradication. Amine formulations are suggested for spring treatments when growth is lush. Esters may be used for fall spraying or if plants are stressed. Fall retreating is critical.

**RESTRICTIONS:** Avoid drift to trees and sensitive broadleaved crops. Do not graze lactating dairy on treated areas for 7 days. Labels for 2,4-D allow harvesting hay 30 days after application and require a 3 day removal period before slaughter.

---

---

## BIENNIAL THISTLES (Musk, Plumeless, Bull, and Scotch Thistle)

---

---

### 2,4-D Ester or Amine

#### 1.5-2 lb ae 2,4-D (1.5-2 lb ae)

**(\$4.35-7.60)**

**Spring.** Apply at rosette stage. May be used in fall or spring; however, other fall treatments are more effective. The low rate has been satisfactory under ideal conditions; 2 lb/A is most consistent. Esters are preferred for pastures; use amines when spraying near trees. Minimum carrier is 10 to 20 gpa. Control is reduced after flower stalks elongate (bolt). Apply when expected high temperature is to exceed 65° F. This treatment is registered for use in grass pasture and range.

**RESTRICTIONS:** Avoid drift to trees and sensitive broadleaf plants. Do not graze lactating dairy on treated areas for 7 days. Labels for 2,4-D allow harvesting hay 30 days after application and require a 3 day removal period before slaughter.

---

### Tordon 22K (picloram)

#### Tordon 22K + 2,4-D (picloram + 2,4-D)

##### .5 pt Tordon 2L (.12 lb ae)

**(\$7.30-11.40)**

##### .38-.5 pt Tordon 2L + 1 lb ae 2,4-D (.1-.12 + 1 lb ae)

##### 2-3 pt Grazon 2.54L (.12-.18 + .5-.75 lb ae)

**Spring or Fall:** Apply at the seedling or rosette stage. Use Tordon at .5 pt for fall; use Tordon at .38 to .5 pt plus 2,4-D at 1 lb ae/A for spring application. Minimum carrier is 10 gpa for ground or 2 gpa for air. Provides excellent control under a wide range of growing conditions. Best fall treatment. Visual effects develop more slowly than for some treatments. Tordon is registered for use in grass pasture and range. Grazon is a premix containing .54 lb picloram + 2 lb 2,4-D amine per gal; 2 pt Grazon is equivalent to .54 pt Tordon + 1 pt 2,4-D 4L.

**RESTRICTIONS:** Do not use in trees. Avoid drift to sensitive broadleaved plants. Avoid contaminating water. Note other label precautions. Treated areas may be grazed. Note 2,4-D restrictions when using tank-mix.  
**Restricted Use Pesticide.**

---

---

## Milestone (*aminopyralid*)

**3-5 oz Milestone 2L (0.05-0.08 lb ae) (\$7.75-12.90)**

**Spring or Fall:** Apply 3-5 oz/A in the spring or summer to plants in the rosette or bolting stages of growth or in the fall to seedlings or rosettes. Apply 4-5 oz/A when plants are in the late bolting through early flowering growth stages. Control after the late bud stage may be improved by tank-mixing 2,4-D at 1 lb ai/A. Use a non-ionic surfactant (NIS) at 0.25-0.5% (1-2 qt per 100 gallons spray solution) under adverse growing conditions, advanced weed growth stages, or pubescent plants.

Milestone may be applied to non-irrigation ditch banks and seasonally dry wetlands, but may not be applied over water or to areas where surface water is present.

**RESTRICTIONS:** There are no grazing restrictions, but aminopyralid may be transferred in manure from livestock that have consumed treated grass within the previous 3 days.

---

## ForeFront (*aminopyralid + 2,4-D*)

**1.5-2 pt ForeFront 3L (0.06-0.08 lb ae aminopyralid + 0.5-0.67 lb ae 2,4-D) (\$10.85-14.45)**

**Spring or Fall:** Apply in the spring to plants in the rosette or bolting stages or in the fall to seedlings or rosettes. Minimum recommended carrier volume is at least 10 gpa for ground application or 2 gpa for aerial application. Use a NIS as recommended for Milestone (0.25-0.5% v/v).

**RESTRICTIONS:** Do not harvest forage for hay within 14 days after application. Aminopyralid residues may be transferred in manure from livestock that have consumed treated forage within the past 3 days. Do not apply more than 2.6 pt/A in a growing season.

---

## Curtail (*clopyralid + 2,4-D*)

**1-2 qt Curtail 2.38L (.1-.19 + .5-1 lb ae) (\$9.30-18.60)**

**Spring:** Apply at rosette to bud stage. Reduced rate of 1 qt/A is frequently used under favorable conditions. Minimum carrier is 10 gpa for ground or 5 gpa for air. Curtail is a premix containing .38 lb clopyralid (Stinger) plus 2 lb 2,4-D amine per gallon. Curtail is labeled for rangeland, grass pasture, CRP grass seedlings, fencerows, and other noncrop areas.

**RESTRICTIONS:** Do not graze dairy cattle in treated areas for 14 days after application. Remove meat animals 7 days before slaughter if grazing treated areas within 2 weeks of application. Do not harvest hay within 30 days after application.

---

## Dicamba Product + 2,4-D (*dicamba + 2,4-D*)

**1 pt dicamba 4L + 1 lb ae 2,4-D (.5-1 lb ae) (\$10.90-11.80)**

**Spring or Fall:** Tank-mix. Apply at the rosette stage. Use dicamba 4L at 1 pt plus 2,4-D at 1 lb ae/A. Rates as low as .5 pt/A dicamba have been successful under ideal conditions. Use the high rate for large rosettes, dense stands, or dry conditions. Use 10 to 20 gpa carrier for ground equipment. May be applied by air using 3 to 10 gpa carrier if there are no sensitive broadleaved crops in the area. This combination is registered for use in grass pasture and range. Refer to Dicamba Product table.

**RESTRICTIONS:** Avoid drift to trees and sensitive broadleaved plants. Do not graze lactating dairy on treated areas for 7 days after application. Labels for 2,4-D allow harvesting hay 30 days after application and require a 3 day removal period before slaughter. Labeling for rates up to 1 pt/A restricts grazing lactating dairy for 7 days or haying for 37 days after application. Note other restrictions for higher rates and note other label precautions.

---

## Overdrive (*diflufenzopyr + dicamba*)

**4-8 oz Overdrive 70DF (.05-.1 + .125-.25 lb ae) (\$10.45-20.90)**

Overdrive is a selective, postemergence herbicide for the control or suppression of many annual, biennial, or perennial broadleaf weeds in pasture, range, and noncrop sites. Overdrive controls weeds by auxin transport inhibition.

## OVERDRIVE (Continued . . . )

**Spring or Fall.** Use rate is 4 to 8 oz/A based on weed species and maturity. Use higher rates for biennial thistles. A maximum of 10 oz can be applied per season in noncropland sites and a maximum of 8 oz/A in pasture, hay, and rangeland sites. Use 1 qt of NIS per 100 gallons of water or MSO at the rate of 1.5 to 2 pt/A. Do not use less than 3 gpa carrier. Rainfast 4 hours after application. Overdrive may be tank-mixed with several labeled tank-mix partners to improve weed control spectrum.

**RESTRICTIONS:** Supplemental label for aerial application in CRP. Do not plant crops for 30 days after last application. Pasture or rangeland treated with Overdrive can be grazed or harvested for livestock feed immediately after application. Do not apply to newly seeded grasses.

---

## **Ally, Cimarron, or Escort (metsulfuron)**

### **Ally or Cimarron + 2,4-D (metsulfuron + 2,4-D)**

**.2-.3 oz Ally 60XP or Cimarron 60DF (.008-.012 lb ai) (\$5.50-39.30)**

**.5-2 oz Escort 60XP (.019-.076 lb ai)**

**.1-2 oz Ally 60XP or Cimarron 60DF + 1-2 pt 2,4-D 3.8L (.004-.008 + .5-1 lb ae)**

**Spring:** Ally is for use in pasture and range. Escort is for western range, right-of-way, and other noncrop sites. Supplemental labeling in 2004 allows .5 oz/A Escort XP for musk thistle at bolt and bud stage to control plants and inhibit seed production. Apply at rosette stage. Control in SDSU tests has been very good with .2 oz tank-mixed with .5 to 1 lb/A 2,4-D. Bolted plants have been controlled in some tests. Cold, dry conditions reduce activity. Rate ranges for Escort in noncrop and western range sites vary depending on the biennial thistle. Musk and plumeless thistle rates range from .5 to 1 oz, while rates for bull and Scotch thistle range from 1 to 2 oz/A. Minimum carrier is 10 gpa for ground or 2 gpa for air; use 5 gpa for Escort above .5 oz. Use NIS at 1 qt/100 gal. Legumes will be injured. Tank-mix .1 to .2 oz metsulfuron product + 1 to 2 pt 2,4-D 3.8L ae per acre to increase control on certain pasture weeds including biennial thistles.

**RESTRICTIONS:** Do not apply on grass grown for seed. No grazing restrictions at listed rate. Hay may be harvested from Escort treated areas. High rates of Escort may suppress grass production. Do not apply to water, lakes, streams, or where there may be runoff into such areas.

---

## **Cimarron Max (metsulfuron + dicamba + 2,4-D)**

**20-5 Acres/.5 oz Part A + 2.5 gal Part B (.01-.038 + .12-.5 + .38-1.5 lb ae) (\$6.70-26.80)**

Cimarron Max herbicide is a twin-pak combination. Part A contains metsulfuron (Ally). Part B contains 1 lb dicamba (Banvel) plus 2.87 lb ae 2,4-D amine per gallon. The use ratio is .5 oz Part A to 2.5 gal of Part B to treat 5 to 20 acres. Refer to following rate table for acres treated. (\$7.50-28.00).

---

<b>Cimarron Max Rate</b>	<b>Part A Rate (oz/A)</b>	<b>Part B Rate (pt/A)</b>	<b>Acres Treated with .5 oz Part A + 2.5 gal Part B</b>
<b>Rate I</b>	<b>0.25</b>	<b>1</b>	<b>20A</b>
<b>Rate II</b>	<b>0.5</b>	<b>2</b>	<b>10A</b>
<b>Rate III</b>	<b>1</b>	<b>4</b>	<b>5A</b>

---

Use Rate I for musk and Scotch thistle and Rate II for plumeless and bull thistle. The Rate II or 10 acre rate provides equivalent of .5 oz Ally 60DF + .5 pt Banvel 4L + 1.5 pt 2,4-D 3.8L per acre. Rates are based on weed species and weeds less than 4 inches tall. Do not apply more than the equivalent of 1.66 oz/A Cimarron Max Part A per year. Add NIS at 2 to 4 pt/100 gal. May be applied by ground or air. No restrictions or waiting period between treatment and grazing for non-lactating animals. Remove meat animals 30 days prior to slaughter. Do not graze lactating dairy within 7 days of treatment. Do not harvest hay for 37 days after treatment.

---

---

**Telar (*chlorsulfuron*)****.5-1.3 oz Telar 75XP (.023-.061 lb ai)****(\$10.25-26.60)**

**Spring:** Apply at rosette stage. Use .5 to 1 oz for musk thistle and 1 to 3 oz/A for bull and Scotch thistle. Minimum carrier is 10 gpa. Add NIS at 1 qt/100 gal. May be tank-mixed with 2,4-D, dicamba, or other labeled tank-mix partners.

**RESTRICTIONS:** There are no grazing restrictions for rates less than 1.3 oz/A. Labeling does not include hay uses. Do not apply to water, such as lakes, streams, or areas where runoff flows into such areas.

---

**Redeem (*triclopyr + clopyralid*)****1.5-2.5 pt Redeem 3L (.42-.7 + .07-.12 lb ae)****(\$17.80-29.70)**

**Spring.** Premix. Apply from rosette to early bolt stage. Add .5 lb/A 2,4-D if weeds are fully bolted. Minimum carrier is 10 gpa for ground or 3 gpa for air. Redeem is labeled for use in range, pasture, CRP grass seedings, fence rows, and noncrop areas. Redeem contains 2.25 lb triclopyr (Garlon) plus .75 lb clopyralid (Stinger) per gallon.

**RESTRICTIONS:** Do not graze or harvest forage for dairy cattle for 14 days after application. Do not harvest hay until the next growing season for dairy or harvest hay for other livestock for 7 days after treatment. Remove meat animals at least 3 days before slaughter. Do not treat banks or bottom of irrigation ditches.

---

**Stinger or Transline (*clopyralid*)****.33-1 pt Stinger 3L or Transline 3L (.12-.38 lb ae)****(\$19.95-60.45)**

**Spring.** Apply from rosette to early bolt stage. Stinger is labeled for use in grass pasture, rangeland, fallow, fence rows, and other noncrop areas. Transline is labeled for noncrop areas, rangeland, and rights-of-way. Transline is marketed only to government and public agencies. Results have been very good in SDSU tests. Use the high rate for late bolt stage. Minimum carrier is 5 gpa; use at least 10 gpa for most ground applications. Stinger or Transline have potential in sites where grass cannot be damaged or where trees limit use of herbicides with harmful soil residual.

**RESTRICTIONS:** Do not contaminate irrigation ditches. No grazing or haying restrictions.

---

---

---

**ABSINTH WORMWOOD (Wormwood Sage)**

---

---

**2,4-D Ester or Amine****2 lb ae 2,4-D (2 lb ae)****(\$5.80-7.60)**

**Spring.** Apply in spring when wormwood is 8 to 10 inches. Rate is 2 lb ae/A. Suggested carrier is 20 gpa. Control is variable. Good coverage and temperature over 65 ° F. improves control. Low-volatile ester formulations are preferred for grass pasture and roadsides. Use amine formulations near trees or where vapor-drift risk is critical for sensitive plants.

**RESTRICTIONS:** Avoid drift to sensitive plants. Do not graze lactating dairy on treated areas for 7 days. Labels allow harvesting hay 30 days after application and require a 3 day removal period before slaughter. Product label restrictions vary.

---

**Milestone (*aminopyralid*)****6-7 oz Milestone 2L (0.09-0.11 lb ae)****(\$15.00-17.50)**

**Spring or Fall:** Apply 6-7 fl oz/A before wormwood is 12 inches tall. May see reduced control with later applications, particularly in drought stressed conditions. Use a non-ionic surfactant (NIS) at 0.25-0.5% (1-2 qt per 100 gallons spray solution) under adverse growing conditions or advanced weed growth stages.

Recommended minimum carrier volume is at least 10 gpa for ground application or at least 3 gpa for aerial application. Greater carrier volumes may improve coverage and control. Removal of old grass litter by mowing or burning may improve coverage and wormwood control. Milestone may be applied to non-irrigation ditch banks and seasonally dry wetlands, but may not be applied over water or to areas where surface water is present.

**RESTRICTIONS:** There are no grazing restrictions, but aminopyralid may be transferred in manure from livestock that have consumed treated grass within the past 3 days.

---

---

**ForeFront (*aminopyralid + 2,4-D*)**

**1.5-2 pt ForeFront 3L (0.06-0.08 lb ae aminopyralid + 0.5-0.67 lb ae 2,4-D) (\$10.85-14.45)**

**Spring or Fall:** Apply before wormwood is 12 inches tall. Recommended minimum carrier volume is at least 10 gpa for ground application or at least 2 gpa for aerial application. Greater carrier volumes may improve coverage and control. Use a NIS as recommended for Milestone (0.25-0.5% v/v).

**RESTRICTIONS:** Do not harvest forage for hay within 7 days after application. Aminopyralid residues may be transferred in manure from livestock that have consumed treated forage within the past 3 days. Do not apply more than 2.6 pt/A in a growing season.

---

**Tordon 22K (*picloram*)****Tordon 22K + 2,4-D Ester (*picloram + 2,4-D*)**

**1 pt Tordon 2L (.25 lb ae) (\$9.60-15.40)**

**.5-1 pt Tordon 2L + 1 lb ae 2,4-D ester (.12-.25 + 1 lb ae)**

**Spring or Fall.** Use Tordon alone or as a tank-mix with 2,4-D ester. Apply in spring before wormwood is over 12 inches. Minimum carrier is 10 gpa for ground or 2 gpa for air. Tordon at 1 pt alone or .5 to 1 pt Tordon plus 1 lb ae/A 2,4-D has provided excellent results in SDSU tests. Results on larger plants have been better than for 2,4-D. Promising as a fall treatment. Tordon is registered for use in grass pasture and range.

**RESTRICTIONS:** Do not use in trees. Avoid drift to sensitive broadleaved crops. Do not graze lactating dairy on treated areas for 7 days. Labels allow harvesting hay 30 days after application and require a 3 day removal period before slaughter. Note other label precautions. There are no restrictions for lactating dairy if Tordon is used alone at this rate. **Restricted Use Pesticide.**

---

---

---

**KNAPWEED SPECIES  
(Russian, Spotted and Diffuse)**

---

**Tordon 22K (*picloram*) *Russian, spotted, and diffuse knapweed***

**1-4 pt Tordon 2L (0.25-1 lb a.e.) (\$11.60-46.25)**

**Spring or Fall.** For diffuse or spotted knapweed, apply 1-2 pt/A in the spring to plants in the rosette or mid-bolting growth stage or in the fall to regrowth. Control at the 1 pt/A rate may be improved by tank mixing with 1 qt/A 2,4-D. For Russian knapweed, apply 2-4 pt/A to actively growing plants in the bud to mid-flowering growth stage or in the fall to regrowth. Some studies have demonstrated very good Russian knapweed control after late-fall applications. Note use restrictions mentioned for other weed species.

---

**Milestone (*aminopyralid*) *Russian, spotted, and diffuse knapweed***

**5-7 oz Milestone 2L (0.06-0.11 lb ae) (\$10.35-18.10)**

**Spring or Fall.** Apply to diffuse or spotted knapweed that is actively growing in the rosette to bolting stage or in the fall. Apply to Russian knapweed in the spring and summer when plants are in the bud to flower growth stage and to dormant plants in the fall. Use a non-ionic surfactant (NIS) at 0.25-0.5% (1-2 qt per 100 gallons spray solution) under adverse growing conditions or advanced weed growth stages. Avoid mowing for 14 days after application to allow for herbicide translocation in the weeds.

Milestone may be applied to non-irrigation ditch banks and seasonally dry wetlands, but may not be applied over water or to areas where surface water is present.

**RESTRICTIONS:** There are no grazing restrictions, but aminopyralid may be transferred in manure from livestock that have consumed treated grass within the previous 3 days.

---

---

**ForeFront (aminopyralid + 2-,4-D) *Russian, spotted and diffuse knapweed***

**2-2.6 pt ForeFront 3L (0.08-0.11 lb ae aminopyralid + 0.67-0.87 lb ae 2,4-D) (\$14.45-18.75)**

***Spring* or *Fall*:** Apply to spotted or diffuse knapweed that is actively growing in the rosette to bolting stage or in the fall prior to a killing frost. Apply to Russian knapweed in the spring or summer when plants are in the early bud to flowering growth stage. Mid-summer or fall applications may not result in immediate visual injury, but control may be observed the following year. Recommended minimum carrier volume is at least 10 gpa for ground application or at least 2 gpa for aerial application. Greater carrier volumes may improve coverage and control. Use a NIS as recommended for Milestone (0.25-0.5% v/v).

***RESTRICTIONS:*** Do not harvest forage for hay within 14 days after application. Aminopyralid residues may be transferred in manure from livestock that have consumed treated forage within the past 3 days. Do not apply more than 2.6 pt/A in a growing season.

---

**2,4-D Ester *Spotted and diffuse knapweed***

**2 lb ae 2,4-D ester (2 lb ae) (\$7.60)**

***Spring.*** Apply at rosette stage. Rate of 2 lb ae/A has provided 95 to 99% control in several SDSU tests. Lower rates may be adequate under ideal conditions. Suggested carrier is 10 to 20 gpa. Apply when expected high temperature is to exceed 65° F.

***RESTRICTIONS:*** Avoid drift to trees and non-target broadleaved plants. Do not graze lactating dairy animals on treated areas for 7 days. Labels allow harvesting hay 30 days after application and require a 3 day removal period before slaughter. Product label restrictions vary.

---

**Curtail (clopyralid + 2,4-D) *Spotted and diffuse knapweed***

**2 qt Curtail 2.38L (.19 + 1 lb ae) (\$18.60)**

***Spring.*** Apply at rosette stage. Minimum carrier is 10 gpa for ground or 5 gpa for air.

Curtail is labeled for rangeland, grass pasture, CRP grass seedings, fencerows, and noncrop areas.

***RESTRICTIONS:*** Do not graze dairy cattle in treated areas for 14 days after application. Remove animals 3 days before slaughter if grazing treated areas within 2 weeks after application. Do not harvest hay within 30 days after application.

---

**Redeem (triclopyr + clopyralid) *Russian, spotted, and diffuse knapweed***

**1.5-4 pt Redeem 3L (.42-.7 + .07-.12 lb ae) (\$17.80-29.70)**

***Spring* or *Fall.*** For spotted and diffuse knapweed, apply 1.5-2.5 pt/A from rosette to early bolt stage. Add .5 lb/A 2,4-D if weeds are fully bolted. For Russian knapweed, apply 3-4 pt/A at the bud to mid-flowering growth stage or to fall regrowth. Some studies have demonstrated very good Russian knapweed control after late fall application. Use the higher recommended rates when weeds are in advanced growth stages or are stressed due to drought or low temperatures. Minimum carrier is 10 gpa for ground or 3 gpa for air. Redeem is labeled for range, pasture, CRP grass seeding, fence rows, and noncrop areas. Redeem contains 2.25 lb triclopyr (Garlon) plus .75 lb clopyralid (Stinger) per gallon.

***RESTRICTIONS:*** Do not graze or harvest forage for dairy cattle for 14 days after application. Do not harvest hay until the next growing season for dairy or harvest hay for other livestock for 7 days after treatment. Remove animals at least 3 days before slaughter. Do not treat banks or bottom of irrigation ditches.

---

---

**Stinger or Transline (*clopyralid*) *Spotted and diffuse knapweed***

**.5-.67 pt Stinger 3L or Transline 3L (.16-.38 lb ae) (\$30.20-40.50)**

**Spring.** Apply from mid bolt to late bud stage. Stinger is labeled for grass pasture, rangeland, fallow, fence rows, and other non-crop areas. Transline is labeled for non-crop areas and rights-of-way. Transline is marketed only to government and public agencies. Use the high rate for most situations. Minimum carrier is 2 gpa; use at least 10 gpa for most ground applications.

Stinger or Transline have potential in sites where grass cannot be damaged or where trees limit use of herbicides with harmful soil residual.

**RESTRICTIONS:** Do not contaminate irrigation ditches. No grazing or haying restrictions.

---

**Plateau (*imazapic*) *Russian knapweed***

**12 oz Plateau 2L (0.18 lb ai) (\$27.25)**

**Fall.** Apply in the fall after leaves begin to die back (senescence). Control may improve as senescence progresses and may still be obtained after full senescence. Apply with MSO at 1 qt/A. Plateau rates greater than 8 oz/A can cause significant grass injury. Do not harvest treated areas for hay until 7 days after application.

---

**Telar (*chlorsulfuron*) *Russian knapweed***

**1-3 oz Telar 75XP (\$20.50-61.50)**

**Spring or Fall:** Apply at the bud to bloom growth stage in early summer or to rosettes in the fall. Registered for use on non-crop, right-of-way, pasture, range, and CRP. However, the maximum rate for pasture/range and CRP is 1.3 oz/A per year. Bluestem, buffalograss, green needlegrass, Indiangrass, and switchgrass may be tolerant to Telar rates up to 0.5 oz/A whereas several wheatgrass varieties, bluegrass, and smooth brome grass may be tolerant to rates over 0.5 oz/A. Minimum carrier is 10 gpa. Add NIS at 0.25% v/v (1 qt per 100 gallons spray solution).

---

---

---

**MULLEIN**

---

---

**Escort or Cimarron (*metsulfuron*)**

**1-2 oz Escort 60XP (.038-.076 lb ai) (\$7.30-39.30)**  
**.1 oz Cimarron 60DF (.004 lb ai)**

**Spring.** Apply at rosette stage. Full coverage is very important. Ally, Cimarron, and Escort labels include pasture and western range uses; Escort also is used in rights-of-way and other noncrop sites. Control with .3 oz in SDSU tests has been marginal; suggest using Escort at .5 to 1 oz/A. Add NIS at 1 qt/100 gal.

**RESTRICTIONS:** Do not apply on grasses grown for seed. No grazing restriction. Allow 3 days after application before harvesting hay if more than 1.66 oz/A Escort is applied. Do not apply to lakes, streams, or sites that may have runoff into such areas. High rates of Escort may suppress grass production.

---

**Telar (*chlorsulfuron*)**

**1-1.3 oz Telar 75XP (.047-.061 lb ai) (\$20.50-26.60)**

**Spring.** Apply at rosette stage. Full coverage is very important for best results. Minimum carrier is 10 gpa. Add NIS at 1 to 2 qt/100 gal. May be mixed with 2,4-D, dicamba, or other labeled tank-mix partners.

**RESTRICTIONS:** There are no grazing restrictions for rates less than 1.3 oz/A. Labeling does not include hay uses. Do not apply to water, such as lakes, streams, or areas where runoff flows into such areas.

---



---

## Milestone (*aminopyralid*)

**7 fl oz Milestone 2L (0.11 lb ae)  
(\$17.50)**

**Spring:** Apply at the rosette stage. Full coverage and use of a surfactant is necessary for best results. Use NIS at 0.25-0.5% v/v (1-2 qt per 100 gallon spray solution). Control has been very good in SDSU trials. Lower Milestone rates have resulted in less control.

**RESTRICTIONS:** There are no grazing restrictions, but aminopyralid may be transferred in manure from livestock that have consumed treated forage within the previous 3 days.

---

## ForeFront (*aminopyralid + 2,4-D*)

**2-2.6 pt ForeFront 3L (0.08-0.11 lb ae aminopyralid + 0.67-0.87 lb ae 2,4-D) (\$14.45-18.75)**

**Spring:** Apply at the rosette stage. Full coverage and use of a surfactant is necessary for best results. Use NIS at 0.25-0.5% v/v (1-2 qt per 100 gallon spray solution). Control has been very good in SDSU trials. Lower ForeFront rates have resulted in less control.

**RESTRICTIONS:** Do not harvest forage for hay within 14 days after application. Aminopyralid residues may be transferred in manure from livestock that have consumed treated forage within the past 3 days. Do not apply more than 2.6 pt/A in a growing season.

---

## Tordon (*picloram*)

### Tordon + 2,4-D Ester (*picloram + 2,4-D*)

**1 qt Tordon 2L (.5 lb ae) (\$15.40-23.20)  
1 pt Tordon 2L + 1 lb ae 2,4-D ester (.25 + 1 lb ae)**

**Spring:** Apply at rosette stage prior to stalk elongation. Use Tordon alone or at a lower rate in a tank-mix with 2,4-D. Treatments have exceeded 90% control in tests in western South Dakota. Surfactant improves penetration through woolly leaf surface. Minimum carrier is 10 gpa for ground and 2 gpa for air. Reduction is apparent for at least 2 years.

**RESTRICTIONS:** Do not use in trees. Avoid drift to sensitive broadleaved crops. Avoid contaminating water. Note other label precautions. Note 2,4-D restrictions when using tank-mix. Restricted Use Pesticide.

---

---

---

## PURPLE LOOSESTRIFE

---

### Aquatic Glyphosate Product (*glyphosate*)

**4 pt aquatic glyphosate 4L (2 lb ae) (\$12.80)**

Aquatic glyphosate is available in several products that are approved for use on aquatic sites. Examples of products include Rodeo, AquaMaster, ClearOut 41 Aquatic, AquaNeat, and AquaStar. Glyphosate is a non-selective, translocated, foliage-applied herbicide, both grasses and broadleaf plants are affected.

**Postemergence.** Apply to actively growing plants at full to late flower. Apply as a broadcast treatment at 4 pt/A or as a spot treatment using hand-held equipment with a 1% aquatic glyphosate solution (1 gal/100 gal or 3 tablespoons/gal) to thoroughly wet foliage. Late summer or fall treatments are best. Apply before killing frost. Rate is minimum required for 65 to 85% control. Use NIS (2 qt/100 gal) with all applications.

This plant is a perennial capable of producing new shoots from buds in the crown areas. Wild types also produce seed. Individual plants can be dug if all small crown pieces are removed. The plant is especially aggressive in wetlands.

Aquatic glyphosate is approved for aquatic uses in lakes, streams, ponds, irrigation ditches, and reservoirs. Allow 7 days before introduction of water into dry ditches. Limit treatment to individual plants if possible.

**RESTRICTIONS:** Do not apply within ½ mile upstream of potable water intake in flowing streams or within ½ mile of intake in standing water. There are no restrictions for irrigation or recreational uses.

---

---

## Habitat (*imazapyr*)

1 pt Habitat 2L (.25 lb ai)

(\$32.15)

Habitat is an aqueous solution to be mixed with water and surfactant for aquatic use and requires spray adjuvants (NIS, MSO, silicone based). Apply 1 pt/A Habitat to actively growing foliage of purple loosestrife in and around standing and flowing water, including estuarine and marine sites. Do not apply more than 6 pt/A per year. Aerial application is restricted to helicopter only.

**RESTRICTIONS:** Application of Habitat can only be made by federal or state government entities or applicators who are licensed or certified applicators making applications under a program sponsored by federal or state government entities. There are no restrictions on livestock consumption of water from the treatment area.

---

## Garlon (*triclopyr*)

.33-3 gal Garlon 3A (.1-9 lb ae)

(\$31.10-282.75)

Garlon 3A may be used within production forests and industrial noncrop sites to control target vegetation in and around standing water sites, such as marshes, wetlands, and banks of ponds and lakes and transition areas between upland and lowland sites. Use enough water to give uniform and complete coverage for best results. NIS is recommended for all foliar applications.

**RESTRICTIONS:** Do not apply directly to un-impounded rivers or streams. Do not apply in ditches or canals used to transport irrigation water. Do not apply where runoff water may flow onto agricultural land. Minimize overspray to open water when making applications to banks or shorelines of moving water sites. There are no restrictions on use of water in treatment area for recreational purposes including swimming and fishing or for livestock consumption.

The maximum rate is 3 gal/A per year for all terrestrial use sites other than range, pasture, forestry sites, and grazed areas. Refer to grazing and haying restriction section of this guide for restrictions.

---

---

## COMMON TANSY

---

## Tordon 22K (*picloram*)

2-3 pt Tordon 2L (.5-.75 lb ae)

(\$23.15-34.70)

**Spring.** Apply before bloom. Results in SDSU tests show 75 to 90% control. The 2 pt/A rate is minimal; adding 1 lb ae/A 2,4-D ester is suggested with the low rate. Apply in minimum of 10 gpa for ground equipment. Tordon is registered for grass pasture and range.

**RESTRICTIONS:** Do not use in trees. Avoid drift to sensitive broadleaved crops. For rates above 1 qt/A, do not harvest hay for 2 weeks or graze dairy animals for 2 weeks and remove slaughter animals 3 days before slaughter. Refer to surface and groundwater precautions. Restricted Use Pesticide.

---

## Escort (*metsulfuron*)

1-2 oz Escort 60XP (.038-.075 lb ai)

(\$19.65-39.35)

**Postemergence.** Apply to actively growing plants. Good spot treatment option. Results have been very good. Add NIS at 1 to 2 qt per 100 gal. Minimum carrier is 10 gpa for ground or 5 gpa for air. May be tank-mixed with 2,4-D or dicamba. Escort may be used in western range, rights-of-way, and other noncrop areas.

**RESTRICTIONS:** Do not apply to lakes, streams, or areas where runoff flows into such areas. No grazing or haying restrictions at this rate.

---

---

## Telar (*metsulfuron*)

1-1.3 oz Telar 75XP (.047-.061 lb ai) (\$20.50-26.60)

**Postemergence.** Apply to actively growing plants. Minimum carrier is 10 gpa for ground. Add NIS at 1 qt/100 gal. May be mixed with 2,4-D, dicamba, or other labeled tank-mix partners.

**RESTRICTIONS:** There are no grazing restrictions for rates less than 1.3 oz/A. Labeling does not include hay uses. Do not apply to water, such as lakes, streams, or areas where runoff flows into such areas.

---

## ST. JOHNSWORT

---

### Tordon (*picloram*)

#### Tordon + 2,4-D Ester (*picloram* + 2,4-D)

1 pt Tordon 2L (.25 lb ae) (\$8.70-15.40)

.5-1 pt Tordon 2L + 1 lb ae 2,4-D (.12-.25 + 1 lb ae)

**Spring or Fall:** Use Tordon alone or as a tank-mix with 2,4-D. Has provided excellent control in SDSU tests. Spring application at bud stage gave 95% control for a 2-year period. Reduce the Tordon rate to .5 pt/A when used with 2,4-D if conditions are favorable. Control has averaged 90 to 95% for one year. Minimum carrier is 10 gpa for ground or 2 gpa for air. Tordon may be used in grass pasture, range, and rights-of-way.

**RESTRICTIONS:** Do not use in trees. Avoid drift to sensitive broadleaved plants. Avoid contaminating water. Note other label precautions. Note 2,4-D restrictions when using tank-mixes. Restricted Use Pesticide.

---

### Milestone (*aminopyralid*)

5-7 fl oz Milestone 2L (0.06-0.11 lb ae) (\$12.50-17.50)

**Spring or Fall:** Apply in the spring at the bud growth stage or in the fall to regrowth. Use NIS at 0.25-0.5% v/v (1-2 qt per 100 gallon spray solution) in adverse growing conditions or to weeds at advanced growth stages.

**RESTRICTIONS:** There are no grazing restrictions, but aminopyralid may be transferred in manure from livestock that have consumed treated forage within the previous 3 days.

---

### ForeFront (*aminopyralid* + 2,4-D)

2-2.6 pt ForeFront 3L (0.08-0.11 lb ae aminopyralid + 0.67-0.87 lb ae 2,4-D) (\$14.45-18.75)

**Spring or Fall:** Apply in the spring at the bud growth stage or in the fall to regrowth. Use NIS at 0.25-0.5% v/v (1-2 qt per 100 gallon spray solution) in adverse growing conditions or to weeds at advanced growth stages.

**RESTRICTIONS:** Do not harvest forage for hay within 14 days after application. Aminopyralid residues may be transferred in manure from livestock that have consumed treated forage within the past 3 days. Do not apply more than 2.6 pt/A in a growing season.

---

### Escort (*metsulfuron*)

.75 oz Escort 60XP (.028 lb ai) (\$14.75)

**Postemergence.** Escort may be used in western range and noncrop areas such as utility sites and rights-of-ways. Apply at the bud to bloom stage; may also be applied to fall regrowth. Results from spot treatments have been excellent. Special 2(ee) labeling for St. Johnswort. Add NIS at 1 qt/100 gal. Minimum carrier is 10 gpa for ground or 4 gpa for air. May be tank-mixed with 2,4-D, dicamba, picloram, triclopyr, and clopyralid.

**RESTRICTIONS:** Do not apply to water, lakes, streams, or areas that may have runoff. No grazing restrictions at this rate.

---

---

## 2,4-D Ester

2-3 lb ae 2,4-D Ester (2-3 lb ae)

(\$7.60-11.40)

**Spring.** Apply in spring at pre-bud stage. Provides suppression; usually 25% reduction from a single application. Repeat treatment required. Amines are not effective. Some grass stunting may be noted with high rates, especially if applied as grass begins forming seedheads. Good coverage is helpful; use 20 gpa carrier.

**RESTRICTIONS:** Avoid drift to trees and sensitive broadleaved plants. Do not graze lactating dairy animals on areas treated for 7 days. Labels allow harvesting hay 30 days after application and require a 3 day removal period before slaughter. Product label restrictions vary.

---

---

## HOUNDSTONGUE

---

---

### Escort XP (*metsulfuron*)

1-2 oz Escort 60XP (.038-.076 lb ai)

(\$19.65-39.35)

**Postemergence.** Escort may be used in western range and noncrop sites. Apply at stalk elongation but before bloom. Results from spot treatments have been very good. Add NIS at 1 qt/100 gal. Minimum carrier is 10 gpa for ground or 5 gpa for air.

**RESTRICTIONS:** Do not apply to water, lakes, streams, or areas that may have run off. No grazing or haying restrictions at this rate.

---

## 2,4-D Ester

2 lb ae 2,4-D ester (2 lb ae)

(\$7.60)

**Spring.** Apply while plants are actively growing but before bloom.

**RESTRICTIONS:** Avoid drift to trees and sensitive broadleaved plants. Do not graze lactating dairy animals on treated areas for 7 days. Labels allow harvesting hay 30 days after application and require a 3 day removal period before slaughter. Product label restrictions vary.

---

### Tordon 22K (*picloram*)

1 qt Tordon 2L (.5 lb ae)

(\$23.15)

**Postemergence.** Apply when plants are actively growing and before flower development.

**RESTRICTIONS:** Do not use in trees. Avoid drift to sensitive broadleaf plants. Avoid contaminating water. Note other label precautions. **Restricted Use Pesticide.**

---

---

## TOADFLAX (Dalmatian and Yellow)

---

---

### **Tordon 22K** (*picloram*)

#### **Tordon 22K + 2,4-D ester** (*picloram + 2,4-D ester*)

**2 qt Tordon 2L (1 lb ae)** **(\$28.85-46.30)**  
**1 qt Tordon 2L + 1.5 lb ae 2,4-D ester (.5+1.5 lb ae)**

**Spring or Fall.** For Dalmatian toadflax, apply to actively growing plants before full bloom or in late summer or fall. Use lower rates for fall application shortly after the first killing frost. Use Tordon in tank-mix with 2,4-D in spring before full bloom. May require annual treatment for 2 to 3 years. Use higher rates (2 qt) for yellow toadflax.

**RESTRICTIONS:** Do not use in trees. Avoid drift to trees or sensitive broadleaf crops. For rates above 1 qt/A, do not harvest for hay within 2 weeks of treatment. Do not graze lactating dairy animals for 2 weeks after treatment. Remove animals 3 days before slaughter if grazing within 2 weeks after spraying. Do not apply into water or wetlands or on inner banks of irrigation or drainage ditches. Risk of leaching is greatest where soils have rapid permeability (such as loamy to sand) and where the underlying aquifer is near the surface. **Restricted Use Pesticide.**

---

### **Telar** (*chlorsulfuron*)

**1-2.5 oz Telar 75XP (.017 + .012 lb ai)** **(\$20.50-51.25)**

**Postemergence.** For dalmatian toadflax, apply 2-3 oz/A and use a high carrier volume (minimum of 24 gallons water/A) if possible. Use NIS at 1 qt/100 gallons. Fall applications may provide more consistent control. For yellow toadflax, apply a minimum of 1.5 oz/A.

**RESTRICTIONS:** Do not allow spray drift onto sensitive crops. There are no grazing restrictions for rates less than 1.3 oz/A. Labeling does not include hay uses. Do not apply to water, such as lakes, streams, or areas where runoff flows into such areas. Special 2(ee) label for these weeds. Note label restrictions.

---

### **Escort** (*metsulfuron*)

**2 oz Escort 60XP (.070 lb ai)** **(\$39.35)**

**Postemergence.** Apply to actively growing plants. Good spot treatment option; spray to wet entire plant. Escort may be used in western range, rights-of-way, and other noncrop areas.

**RESTRICTIONS:** Do not apply to lakes, streams, or areas where runoff flows into such areas. No grazing restrictions at this rate. Allow 3 days after application before harvesting hay if more than 1.66 oz/A Escort is applied. Special 2(ee) label for these weeds.

---

### **Plateau** (*imazapic*)

**8-12 oz Plateau 2L (.12-.18 lb ai)** **(\$18.20-27.25)**

**Fall.** Label recommends 12 oz/A of Plateau plus 2 pt/A MSO for control of dalmatian toadflax. For best results apply to the basal growth in the fall, after the first hard frost. The plant can have the top 25% showing necrotic tissue; however, there should be green stem and leaf tissue remaining. Applications made prior to this timing will result in poor control. Grass tolerance is an important issue.

**RESTRICTIONS:** Treated areas may be grazed. Do not harvest for 7 days after treatment. Plateau is being marketed only through public agencies and organizations. However, it is labeled and may be applied to private as well as public lands.

---

---

---

---

## PUNCTUREVINE

---

---

### 2,4-D Amine or Ester

2 lb ae 2,4-D (2 lb ae) (\$3.80-7.60)

**Postemergence.** Apply when new seedlings appear. Tank-mixes with non-selective herbicides in noncrop areas will improve year-long control. Apply in 10 to 20 gpa carrier.

**RESTRICTIONS:** Note 2,4-D label restrictions. Avoid drift to non-target plants.

---

### Telar (*chlorsulfuron*)

1.3 oz Telar 75XP (.061 lb ai) (\$26.60)

**Postemergence.** Apply in the late fall or very early spring before spring growth. Moisture is required to activate.

May be used with other selective or bareground herbicides.

**RESTRICTIONS:** Avoid drift to sensitive crops. There are no grazing restrictions for rates less than 1.3 oz/A. Labeling does not include hay uses. Do not apply to water, such as lakes, streams, or areas where runoff flows into such areas.

---

---

## POISON HEMLOCK

---

---

### 2,4-D Ester + Dicamba Product (2,4-D + dicamba)

1 lb ae 2,4-D Ester + .5 to 1 pt dicamba 4L (1 + .25-.5 lb ae) (\$7.80-11.80)

**Fall or Early Spring.** Apply at fall rosette stage or to new growth in early spring. Use 10 to 20 gpa carrier for ground equipment. May be applied by air using 3 to 10 gpa carrier if there are no sensitive broadleaved crops in the area. Refer to Dicamba Product table.

**RESTRICTIONS:** Avoid drift to trees and sensitive broadleaved plants. Do not graze lactating dairy on treated areas for 7 days after application. Labels for 2,4-D allow harvesting hay 30 days after application and require a 3 day removal period before slaughter. Labeling for rates up to 1 pt/A restricts grazing lactating dairy for 7 days or haying for 37 days after application.

---

### Tordon 22K + 2,4-D (*picloram* + 2,4-D)

.38 - .5 pt Tordon 2L + 1 lb ae 2,4-D (.1-.12 + 1 lb ae) (\$7.60-15.20)

2-4 pt Grazon 2.54L (.12-.24 + .5-1 lb ae)

**Spring.** Apply in early spring before flower stalks fully elongate. Minimum carrier is 10 gpa for ground or 2 gpa for air. Tordon is registered for use in grass pasture and range. Grazon is a premix containing .54 lb picloram + 2 lb 2,4-D amine per gal; 2 pt Grazon is equivalent to .54 pt Tordon + 1 pt 2,4-D 4L.

**RESTRICTIONS:** Do not use in trees. Avoid drift to sensitive broadleaved plants. Avoid contaminating water. Note other label precautions. Treated areas may be grazed. Note 2,4-D restrictions when using tank-mix.  
**Restricted Use Pesticide.**

---

---

## SALT CEDAR (*Tamarix species*)

---

### Arsenal, Stalker, Habitat, or Polaris (*imazapyr*)

#### 4-6 pt Arsenal 2L or Polaris 2L (1-1.5 lb ai)

(\$144.40-216.50)

**Foliar:** Apply Arsenal at 2 qt + 0.25% v/v NIS or 1 qt/A MSO or spot treat with a 1% solution of Arsenal + 0.25% v/v NIS or 0.25% v/v MSO. Apply to actively growing foliage. When cambium is brown (about 2 years) burn off dead brush and replant to desirable species.

**Cut Stump:** Apply a solution of Arsenal at 12 oz in 1 gal of water + .25% v/v MSO to the cambium area of freshly cut stump surfaces or to fresh cuts on the stem. Applications can be made anytime of the year except during periods of heavy sap flow in spring. Do not over apply solution causing run off or puddling.

**Cut Stubble:** Apply Arsenal within 2 weeks after mowing or cutting brush. To suppress or control resprouting, uniformly apply 1 to 2 pt/A to cut area. Arsenal may be tank-mixed with Tordon K or 22K to aid in control. The addition of 5% v/v penetrating agent may aid uptake. Direct applications can increase potential root uptake of desirable trees causing injury or death.

**Frill or Girdle:** Make cuts through the bark with a hatchet, machete, etc. at about equal intervals around the tree. Make one cut for every 3 inch of DBH (diameter at breast height) on targeted tree. Spray or brush a solution of 12 oz/gal water + .25% v/v MSO into each cut until thoroughly wet.

#### 1-2 pt Stalker 2EC (.25-.5 lb ai)

(\$48.30-96.50)

Stalker is an emulsifiable concentrate that can be mixed with water, diesel oil, recommended seed oils, and penetrating oils. Mix 8 to 12 oz Stalker with one gallon water, diesel oil, or penetrating oil. May be applied as a spray to cut stump and frilling cuts, when mixed with water or penetrating oil. Basal treatments should be mixed with penetrating oil. Cut stubble treatments need to be applied within 2 weeks after mowing or cutting.

#### 3 qt Habitat 2L (1.5 lb ai)

(\$192.95)

Habitat is an aqueous solution to be mixed with water and surfactant approved for aquatic use. For aerial application (helicopter only), apply 2 qt/A Habitat + 0.25% v/v NIS to actively growing foliage during flowering. For spot spraying use 1% solution of Habitat + 0.25% v/v NIS and spray to wet foliage. Wait at least 2 years after application before disturbing treated saltcedar. Earlier disturbance can reduce overall control. Habitat can also be applied as cut stump, cut stem, and frill or girdle treatments.

**RESTRICTIONS:** Use precautions to minimize or eliminate spray drift. Be sure appropriate buffer zones can be maintained to prevent spray drift out of target area. Special local need labeling (Sec 24(c)) for South Dakota allows grazing treated areas. Remove livestock for slaughter from treated areas for 30 days following treatment. Do not harvest for hay for 30 days after treatment. Follow all label restrictions that apply to aquatic sites.

---

### Garlon 4 or Garlon 4 Ultra (*triclopyr*)

#### 1-8 qt Garlon 4L (1-8 lb ae)

(\$23.50-188.50)

Garlon is recommended for the control of unwanted woody plants and broadleaf weeds. Do not exceed 8 qt/A per year. Basal bark or cut stump treatments are most effective.

**Cut Stump:** Apply anytime the herbicide does not freeze and the tree is not frozen. Cut stumps should be treated immediately (less than 1 hour) after cutting. The outer 2 inches of the stump should be sprayed. Coverage is essential for root kill. Use a 50% solution of 2 qt Garlon 4L + 2 qt water or a 25 to 33% solution of 1 qt Garlon 4A + 3 qt oil or 1.3 qt Garlon 4A + 2.7 qt oil for a 1 gallon mix. Approved oils include basal oils, diesel, kerosene, or MSO.

**Basal Bark:** Use a sprayer with the nozzle adjusted to deliver a narrow, cone shaped spray. Apply the Garlon solution lightly but evenly on the plant stem or trunk up to 12 to 15 inches from the ground. Apply to all sides of every stem. Do not apply to the point where runoff causes puddling at the crown or root collar. Older plants with rougher bark may require stems to be treated higher (15 to 18 inches).

**Foliar:** Use Garlon 4L at rates of 1 to 8 qt/A. A surfactant is allowed but must be mixed in the solution before Garlon. High and low volume foliar treatments are labeled when applied with ground equipment.

**RESTRICTIONS:** Refer to grazing and haying restriction section for precautions and restrictions.

---

## SHELTERBELTS (Does not include fruit trees)

---

### Glyphosate Products (*Numerous*)

(\$19.40-44.90)

**2-4 qt glyphosate 3 lb ae (1.5-3 lb ae)**

**1.6-3.2 qt 3.7 lb ae (1.5-3 lb ae)**

**1.5-3 qt 4 lb ae (1.5-3 lb ae)**

**1.4-2.8 qt 4.5 lb ae (1.5-3 lb ae)**

**2.3-4.6 lb 65DS ae (1.5-3 lb ae)**

**Postemergence.** Apply when weeds are actively growing and at boot or bud to bloom stage. Fall application is more effective than spring. Canada thistle is reduced 75 to 85%; field bindweed 50 to 75%. Field bindweed control is more variable. Apply in 10 to 40 gpa carrier. Rates are 3 qt for Canada thistle, and 4 qt of 3L ae/A for field bindweed. Adjust rates for other formulations.

Products contain different salts of glyphosate acid, including several "Roundup" products, "Touchdown" products, "Glyphomax" products, Glyphos and others. Rates for formulations are listed according to acid equivalent content. Check specific product labeling. Roundup Pro or Touchdown Pro labeling includes tree plantings and noncrop sites. Other products limit use to trees and noncrop areas associated with agricultural sites. Product concentration also varies. Glyphosate is a nonselective, foliar, translocated herbicide. There is no soil residual activity.

**RESTRICTIONS:** Avoid spray or drift contact on green leaves, stems, or new bark. Note other label precautions.

---

### 2,4-D Amine

**1-1.5 lb ae 2,4-D (1-1.5 lb ae)**

(\$2.90-4.40)

**Postemergence.** Selective, translocated herbicide for broadleaved weeds. This herbicide is useful to reduce stands of perennial weeds including field bindweed, Canada thistle, or leafy spurge. Labeling is for noncrop areas; tree uses are not listed. Apply when weeds are actively growing and at bud stage. Requires retreatment in fall. Apply 1 to 1.5 lb ae 2,4-D amine/A. Suggested carrier is 40 gpa. Use no more than 20 psi pressure to produce coarse droplets and reduce risk of drift. Spray when it's calm and expected high temperature is below 75° F.

**RESTRICTIONS:** Very small amounts of herbicide from vapor or droplet drift can seriously damage or kill deciduous trees. Some leaf burn may be noted. Conifers are somewhat less sensitive, especially when not actively growing. Avoid heavy application over tree root zone. Suggested use is limited to special situations where risk of exposure to trees can be assumed. Labelers will not be responsible for damage to trees.

---

### Plateau (*imazapic*)

**8-12 oz Plateau 2L (.12-.18 lb ai)**

(\$18.20-27.25)

**Preemergence and Postemergence.** Labeled for use in approved brush and tree species. Not intended for use on nursery, orchard, ornamental plantings, new plantings, or seedling trees. Plateau controls mustard, smartweed, crabgrass, foxtail, pigweed, lambsquarters, leafy spurge, woodsorrel, foxtail barley, bedstraw, and several other grasses and broadleaved weeds. Plateau is being marketed only through public agencies and organizations. However, it is labeled and may be applied to private as well as public lands.

Early postemergence suggested for most weeds; however there is residual activity for preemergence effect. Always add (MSO) at 1.5 to 2 pt when using less than 30 gpa carrier. For carrier over 30 gpa, use MSO or COC at 1 gal/100 gal. AMS may also be added. NIS may be used in place of seed oil in some situations.

Apply Plateau as a directed spray below the foliage for best selectivity. Some chlorosis may be noted. Suggest use on a limited basis to determine tolerance. Labeled species for directed application include green ash, boxelder, red cedar, cottonwood, hackberry, juniper, locust, sugar maple, oak, white pine, serviceberry, and walnut. Plateau may be mixed with Pendulum or other herbicides approved for use.



---

## Stinger (*clopyralid*)

**.25-.66 pt Stinger 3L (.2-.5 lb ae)**

**(\$15.10-39.90)**

**Postemergence.** Labeled for over-the-top application on certain species of Christmas tree plantings, including Douglas, Fraser, grand, balsam and noble fir; blue spruce; and lodgepole, Ponderosa, and white pine. Useful to control emerged broadleaves such as Canada thistle, knapweeds, and some annuals. Very effective on Canada thistle. Use the high rate for perennials. Maximum for blue spruce is .5 pt/A. Reports indicate good tree tolerance. Do not apply to first-year transplants. Do not use additives.

---

## NOXIOUS WEED CONTROL in CULTIVATED CROPLAND (Fallow, After Harvest)

Fallow and after-harvest periods when crops are not planted offer an opportunity to use special herbicides which will substantially reduce stands of Canada thistle, field bindweed, and other perennials. Results are best if weeds are growing actively and have 6 to 8 inches of growth. Early bud stage is best for most weeds.

Late fall applications are usually more effective than those applied in June or July, if fall conditions favor good growth. Discontinue tillage or herbicide use so perennials will have adequate growth when treated. Blade or sweep tillage is more effective than herbicide if it is dry or if fall regrowth is limited. Cultivation 10 to 14 days after spraying usually provides additional stand reduction. Apply herbicides before weeds are damaged by frost; however, light frost before spraying is not harmful.

Perennial control is often variable. Variability increases as rates are reduced. Use caution to prevent drift to adjacent crops.

---

## 2,4-D

**1-3 lb ae 2,4-D (1-3 lb ae)**

**(\$2.90-11.40)**

Amine or ester forms are available. Use 1.5 lb ae/A of ester for leafy spurge; 1 lb ae/A of amine or ester for field bindweed and Canada thistle. Amine is preferred in spring or early summer when growing conditions are good. Use ester forms for less favorable conditions. Usually provides 10 to 25% stand reduction. Experimental tests using high rate (2 to 3 lb ae/A) have provided improved performance when applied in late September or October. Treated areas may be tilled 2 weeks after application. No restrictions for spring planted crops after fall application. A 15-day interval between application and planting winter grain is suggested.

---

## Dicamba Products (*dicamba*)

**2-4 pt dicamba 4L (1-2 lb ae)**

**(\$16.00-32.00)**

Dicamba at 2 to 4 pt/A is very effective on Canada thistle. It is less effective on leafy spurge. Late fall treatments give very good field bindweed control; mid-summer treatments have not been consistently better than with 2,4-D. Corn or sorghum may be planted the following year. Soil residual limits late fall use in some rotations. Allow 45 days during which soil is not frozen for each pint before planting wheat, oats, or barley. Sunflowers and legumes are not labeled for planting the next season. May be used in grass pasture and fence lines. Refer to Dicamba Products table.

---

## Dicamba Products + 2,4-D (*dicamba + 2,4-D*)

**1-2 pt dicamba 4L + 1 lb ae 2,4-D (.5-1 + 1 lb ae)**

**(\$10.90-19.80)**

Tank-mix. Label rates are 1 to 4 pt/A dicamba plus 2,4-D amine or ester (suggested 1 lb ae/A). Primarily for Canada thistle where dicamba rate must be reduced. Combinations using less than 1 pt/A may be slightly better than 2,4-D alone but are less consistent than higher rates. Follow crop rotation guidelines as for dicamba alone. Labeling includes fallow or stubble to be planted to wheat. Lower rates give suppression; higher rates give seasonal control.

---

---

**Glyphosate Products (Numerous) (\$19.50-44.90)****Glyphosate Products + 2,4-D (glyphosate + 2,4-D)****Glyphosate Products + Dicamba Products (glyphosate + dicamba)**

There are several products containing glyphosate including "Roundup" products, "Touchdown" products, "Glyphomax" products, Glyphos, and others. Refer to specific product for surfactant or ammonium sulfate additive requirements. No restriction on labeled rotational crops.

**2-4 qt 3 lb ae (1.5-3 lb ae)**

**1.6-3.2 qt 3.7 lb ae (1.5-3 lb ae)**

**1.5-3 qt 4 lb ae (1.5-3 lb ae)**

**1.5-3 qt 4.17 lb ae (1.5-3 lb ae)**

**1.4-2.8 qt 4.5 lb ae (1.5-3 lb ae)**

**2.3-4.6 lb 65DS (1.5-3 lb ae)**

All product rates listed are for 3L ae product; refer to the Glyphosate Table for equivalent product rates for other formulations. Minimum label rate is 4 qt for field bindweed or 2 qt/A for Canada thistle. Full rates give better control in mid-summer; however, in late fall 2 qt often appears similar to 4 qt/A. Canada thistle control is very good (75 to 95%). Primarily for patch treatment. Weeds should be actively growing and at bud stage. Do not cultivate in the fall or spring before application. Fall application preferred, especially for field bindweed. Lower rates give more variable results. Rate of 1 qt/A suppresses Canada thistle; 1 pt/A plus 2,4-D or dicamba also suppresses several perennials. Very low rates intended for annual weeds frequently give topgrowth burn but do not consistently reduce stands.

**1 pt glyphosate 3 lb ae + .5 lb ae 2,4-D (.38+.5 lb ae) (\$4.00-10.25)**

Intended for suppression in annual control program. Tank-mix. Use 1 pt glyphosate 3L ae plus .5 lb 2,4-D ae/A for field bindweed. All product rates for glyphosate are for 3L ae products; refer to the Glyphosate Table for equivalent product rates. Refer to specific product for surfactant requirements. The addition of ammonium sulfate is suggested.

**1 pt glyphosate 3 lb ae + .5 pt dicamba 4L (.38-.25 lb ae) (\$7.00)**

Intended for field bindweed and Canada thistle suppression in an annual control program. Tank-mix. Fallowmaster commercial premix does not include perennial suppression labeling. Refer to label guidelines.

---

**Tordon 22K (picloram)****Tordon 22K + 2,4-D (picloram + 2,4-D)**

Federal labeling includes use in fallow cropland at low rates alone or with 2,4-D as a broadcast application or at high rates as a spot treatment. Use only on land to be planted to grass, barley, oats, wheat, flax or fallowed. Allow a minimum of 45 days with soil temperature over 40° F. between application of .5 pt Tordon; increase to 90 days if .5 to 1 pt is used. Sensitive broadleaved crops should not be planted for 36 months. Spot treatment rates of 1 to 4 qt/A will injure crops for two years. Amine form of 2,4-D preferred for thistles and bindweed growing actively. Use ester form for leafy spurge, hoary cress, Russian knapweed, or for late fall spraying.

**1-2 qt Tordon 2L (.5-1 lb ae) (\$23.15-46.30)**

**SPOT TREATMENT in CULTIVATED FIELDS:** Intended for small patches where crop damage can be tolerated for two seasons. Effective for Canada thistle, field bindweed, and leafy spurge. Plan to retreat area for regrowth in the second or third season. Rates over 1 pt/A should not exceed 10% of the immediate field in any one year. Rotate to crops as for broadcast application. **Restricted Use Pesticide.**

**.5-1 pt Tordon 2L + .5-1 lb ae 2,4-D (.12-.25 + .5-1 lb ae) (\$7.20-14.40)**

**BROADCAST FALLOW for SMALL GRAIN:** For field bindweed and perennial thistle, use .5 to 1 pt Tordon plus .5 to 1 lb 2,4-D ae/A after grain harvest. A second application may be made the next season during fallow using the low Tordon rate. Between small grain crops .5 pt Tordon plus .5 to 1 lb ae 2,4-D may be used after harvest as an annual control program. Field bindweed control has been very good in SDSU tests. The lower Tordon rate with the high 2,4-D rate reduces risk of crop damage from soil residual. (Effects on spring grain have been noted in some tests). **Restricted Use Pesticide.**

---

## **Curtail** (*clopyralid + 2,4-D*)

**4 pt Curtail 2.38L (.19-1 lb ae)**

**(\$18.60)**

Curtail is intended for Canada thistle. Curtail is a premix containing .38 lb clopyralid + 2 lb 2,4-D per gallon. Labeled for fallow cropland. Canada thistle control has been very good. Stand reduction the following season is fair to good, considerably better than 2,4-D alone. The full rate gives 70 to 75% reduction in favorable conditions; 2 pt gives good seasonal suppression and some stand reduction. Apply before bud stage. Do not cultivate for 2 to 3 weeks. Crop rotation the following season is limited to corn, wheat, barley, oats, or grasses. Labeled crops should not be planted for 30 days after application.

---

## **AQUATIC**

---

### **Aquatic Glyphosate Products** (*glyphosate*)

**3 to 7.5 pt Aquatic Glyphosate 4L (1.1-2.8 lb ae)**

**(\$9.70-24.00)**

Aquatic glyphosate is available in several products that are approved for use on aquatic sites. Examples of products include Rodeo, Aquamaster, ClearOut 41, AquaNeat, and AquaStar. Most aquatic glyphosate products are formulated as a liquid containing 4 lbs glyphosate acid per gallon or 5.4 lb glyphosate, isopropylamine salt per gallon.

Labeled for aquatic and noncrop sites, including lakes, rivers, ponds, reservoirs, drainage ditches, and similar sites. Glyphosate is nonselective, both grasses and broadleaves are affected. There is no soil residual activity.

Aquatic glyphosate controls several species of brush, perennial or annual grasses, and broadleaves. Primary use is to control Canada thistle and other perennials in aquatic sites where other treatments cannot be used.

Rates are 3 to 4.5 pt for Canada thistle or 6 to 7.5 pt/A for field bindweed. Use a 1.5% solution (4 T/gal) for hand-held equipment. Add 2 qt/100 gal solution of labeled surfactant. Apply in 3 to 20 gpa carrier. Rates for other aquatic and noncrop site weeds are based on specific weeds species on the label.

**RESTRICTIONS:** Do not apply within ½ mile upstream of a potable water intake, in moving water, or within a half mile of a potable water intake in streams, ponds, or reservoirs. Allow a minimum of 7 days after treatment before reintroducing water if applying in dry ditches. There is no restriction on the use of treated water for irrigation, recreation, or domestic purposes.

---

## **2,4-D**

**1-2 lb ae 2,4-D (1-2 lb ae)**

**(\$2.90-7.60)**

Certain amine and ester products are labeled for aquatic sites, including areas around marshes, ponds, irrigation ditches, streams, and lakes. Some products are labeled for aerial application in aquatic sites. Check labels for correct labeling. Do not apply to more than 1/3 to 1/2 of a lake or pond in any one month because excessive decaying vegetation may deplete oxygen control in water and kill fish. Do not contaminate water used for irrigation or domestic purposes. Useful for controlling Canada thistle, perennial sow thistle, leafy spurge, or other broadleaf weeds.

**RESTRICTIONS:** Refer to label for application guidelines.

---

## **Habitat** (*imazapyr*)

**1-6 pt Habitat 2L (.25-1.5 lb ai)**

**(\$32.15-192.95)**

Habitat is an aqueous solution to be mixed with water and surfactant approved for aquatic use. Applications of 1 to 6 pt/A may only be made to control undesirable emergent and floating aquatic vegetation in or around standing and flowing water, including estuarine and marine sites. Rates are based on aquatic weed species on the label. Do not apply more than 3 qt/A per year. Applications may be made to private waters that are still, such as ponds, lakes, and drainage ditches where there is minimal or no outflow to public waters. Applications may be made to public waters such as ponds, lakes, reservoirs, marshes, drainage ditches, canals, streams, rivers, and other slow-moving or quiescent bodies of water. Applications to public waters can only be made by state or federal agencies or applicators licensed or certified as aquatic pest control applicators.

## HABITAT (Continued . . . )

**PRECAUTIONS or RESTRICTIONS:** Do not apply directly to water within ½ mile upstream of an active potable water intake in flowing water or within ½ mile of an active potable water intake in a standing body of water such as lakes, ponds, or reservoirs. There are no restrictions on livestock consumption of water from a treated area. Special local need labeling (Sec 24(c)) for South Dakota allows grazing treated areas. Slaughter livestock must be removed from treated areas for 30 days following treatment. Do not cut treated areas for hay for at least 30 days after treatment.

---

### Garlon 3A (*triclopyr*)

**.33-3 gal Garlon 3A (.1-9 lb ae)**

**(\$24.00-218.30)**

Garlon 3A may be used within production forests and industrial noncrop sites to control target vegetation in and around standing water sites, such as marshes, wetlands, and banks of ponds and lakes and transition areas between upland and lowland sites. Rates are based on broadleaf and woody plant species on the label.

**PRECAUTIONS and RESTRICTIONS:** Do not apply directly to un-impounded rivers or streams. Do not apply on ditches or canals used to transport irrigation water. Do not apply where runoff water may flow onto agricultural land. Minimize overspray to open water when making applications to banks or shorelines of moving water sites. There are no restrictions on use of water in treatment area for recreational purposes including swimming and fishing or for livestock consumption.

The maximum rate is 3 gal/A per year for all terrestrial use sites other than range, pasture, forestry sites, and grazed areas. Refer to grazing and haying restriction section of this guide for restrictions.

---

## NONCROP and SPECIAL SITUATIONS

Includes treatments for special situations. Several treatments described in other sections can also be used on noncrop areas.

---

### Arsenal (*imazapyr*)

**1-6 pt Arsenal 2L (.25-1.5 lb ai)**

**(\$36.10-216.60)**

Arsenal is registered for noncrop areas. Arsenal is absorbed through the root or leaves and is translocated in the plant. Postemergence applications are preferred for most situations. It is especially effective for field bindweed (1 pt) but also controls dock, bull thistle, and poison ivy (3-4 pt) and Canada thistle, sow thistle, milkweed, and many brush species (4-6 pt/A) in 1 gallon of water. Use 30 to 60 gpa carrier plus NIS. Results have been very good.

**RESTRICTIONS:** It is not approved for use in cultivated fields, pasture, or hayland. It is not to be used on sidewalks, driveways, and similar areas. Avoid use where the herbicide may come in contact with roots of desirable trees and shrubs.

---

### Krenite (*fosamine*)

**1.5-3 gal Krenite 4L (6-12 lb ai)**

**(\$78.90-157.80)**

Labeled for noncrop areas. It may be used adjacent to and in areas surrounding domestic water reservoirs, streams, ponds, and rights-of-way. Labeling does not include cultivated fields or areas that are hayed or grazed.

Primarily used as a foliar spray to control brush. Treated portions of brush are killed. Also controls and suppresses leafy spurge. It is an option where aquatic limitations preclude the use of other treatments.

Apply as a spray between full bloom and seed set. Has provided 40 to 60% stand reduction in some tests. Follow-up treatments are required for greater permanent reduction. Underground shoots are inhibited and existing topgrowth is controlled. The higher rates are suggested for best results. Mix the desired amount of herbicide in 100 gal of water and thoroughly wet foliage. For small areas, use a 3% solution (8 T/gallon) and spray until foliage is wet. Add crop oil at 1 qt/100 gal of solution. Krenite is formulated as a liquid containing 4 lb/gal.

**RESTRICTIONS:** Do not use in food crops. Avoid spray contact with woody plants and other desirable plants.

---

## Paramount (*quinclorac*)

8 oz Paramount 75DF (.38 lb ai)

(\$25.90)

**Spring.** Primarily for field bindweed. Also suppresses leafy spurge and perennial thistle. Controls annual grass but does not cause permanent damage to established grass. Apply 8 oz/A for most perennials. Has excellent activity on field bindweed. Paramount is labeled for use in roadsides, fence lines, rights-of-way, and other noncrop areas and fallow. Apply with ground equipment. Add 2 pt COC or MSO plus .5 to 1 gal 28% N or 2.5 lb AMS per acre. Apply to active weed growth.

**RESTRICTIONS:** Grass in treated areas cannot be grazed or harvested for forage. Follow crop rotation restrictions for fallow application.

---

## Glyphosate Pro Products (*glyphosate*)

(\$4.25-20.25)

**1-5 qt 3 lb ae (.75-3.75 lb ae)**

**.8-4 qt 4 lb ae (.75-3.75 lb ae)**

**1.2-5.9 lb dry (.75-3.75 lb ae)**

Glyphosate Pro is available in several products having different formulations and concentrations. Examples of products with 3 lb acid equivalent (4 lb active ingredient) include Roundup Pro, Touchdown Pro, GlyStar Pro, ClearOut Pro Plus, GlyPro Plus, and others. Examples of products that contain 4 lb acid equivalent (5.4 lb active ingredient) include Roundup Pro Concentrate, GlyPro, and others. There are also dry products, Roundup Pro Dry (71.4% act; 64.9% ae) is an example.

**Postemergence:** Foliar, non-selective, translocated herbicide. No soil residual. Weeds should be actively growing. Controls annuals and perennials. Useful in sites where risk of movement or root injury to trees or shrubs limits use of long-term residual products. Low rates are for annuals; high rate for perennials. Use on weeds like tansy or St. Johnswort at bud or bloom stage in noncrop sites. Results in SDSU tests show 80 to 90% control of tansy. Retreatment is required in the fall of the following year.

Labeling lists many specific noncrop sites including parks, airports, ornamental nurseries, sod farms, trim and edge, bareground, cut stumps, roadsides, guardrails, schools, apartments, and substations. Agriculture labeled glyphosate products contain noncrop labeling only as it pertains to agriculture sites.

**RESTRICTIONS:** Avoid contact with foliage, green stems, etc. of desirable plants and trees as severe injury or destruction may result. Avoid drift when applying near desirable plants and crops.

---

---

## 2,4-D LABEL RESTRICTIONS and NONCROP LABELING

PRODUCT: SD Registration or by Labeler	EPA Reg. #	2,4-D Acid Equiv.	Lbs. acid/gal	Non- crop	Right- of- Way	Aerial Application	Aquatic
<b><u>Agriance L.L.C. (1381)</u></b>							
Agriolutions 2,4-D Amine 4	1381-103	39.3%	3.8	✓	✓	✓	Y*
Agriolutions 2,4-D LV4	1381-102	44.0%	3.8	✓	✓	✓	N
Agriolutions 2,4-D LV6	1381-101	59.1%	5.6	✓	✓	✓	N
Agriolutions Phenoxo 088	42750-36-9779	30.8%	2.8	✓	✓	✓	N
Agriolutions SWB 2,4-D LV4	42750-22-9779	40.9%	3.76	✓	✓	✓	N
<b><u>Albaugh (42750)</u></b>							
AgriStar 2,4-D Amine 4	42750-19	38.9%	3.8	✓	✓	✓	Y*
AgriStar 2,4-D LV4	42750-15	42.5%	3.8	✓	✓	✓	N
AgriStar 2,4-D LV6	42750-20	57.4%	5.5	✓	✓	✓	N
AgriStar Solve 2,4-D	42750-22	40.9%	3.76	✓	✓	✓	N
AgriStar D-638	42750-36	30.8%	2.8	✓	✓	✓	N
<b><u>Helena Chemical Co. (5905)</u></b>							
2,4-D amine 4	42750-19-5905	38.9%	3.8	✓	✓	✓	Y*
2,4-D LV6	42750-20-5905	57.4%	5.5	✓	✓	✓	N
2,4-D LV4	42750-15-5905	42.5%	3.8	✓	✓	✓	N
Barrage H.F. (ester)	5905-529	51.8%	4.7	✓	✓	✓	N
Opti-Amine	5905-501	38.8%	3.8	✓	✓	✓	Y*
Hardball	5905-549	19.6%	1.7	✓	✓	✓	Y*
Helena	5905-542	19.6%	1.7	✓	✓	✓	Y*
<b><u>NuFarm America's &amp; NuFarm Turf and Specialty (71368)</u></b>							
Esteron 99 Concentrate	62719-9-71368	44.1%	3.8	✓	✓	✓	N
Weedar 64	71368-1	38.9%	3.8	✓	✓	✓	Y*
Weedone 638	71368-3	30.8%	2.8	✓	✓	✓	N
Weedone 650 Solventless	35935-6-71368	88.8%	5.6	✓	✓	✓	N
Weedone LV4 EC	228-139-71368	44.6%	3.8	✓	✓	✓	N
Weedone LV4 Solventless	71368-14	41.5%	3.8	✓	✓	✓	N
Weedone LV6 EC	71368-11	57.5%	5.4	✓	✓	✓	N
Riverdale Weeddestroy AM-40 Amine Salt	228-145	39.3%	3.8	✓	✓	✓	Y
Riverdale Formula 40	228-357	36.5%	3.67	✓	✓	✓	N
<b><u>PBI Gordon (2217)</u></b>							
Hi-Dep Broadleaf Herbicide	2217-703	38.6%	3.8	✓	✓	✓	N
<b><u>TenKoz (55467)</u></b>							
Tenkoz 638	42750-36-55467	30.8%	2.8	✓	✓	✓	N
Tenkoz 2,4-D Amine 4	42750-19-55467	38.9%	3.8	✓	✓	✓	Y*
Lo-Vol 4 2,4-D Low Volatile	42750-15-55467	42.5%	3.8	✓	✓	✓	N
Lo-Vol 6 2,4-D Low Volatile	42750-20-55467	57.5%	5.5	✓	✓	✓	N
<b><u>UAP/Loveland Chemical (34704)</u></b>							
Clean Amine 4 2,4-D	34704-120	38.6%	3.74	✓	✓	✓	Y*
Low Vol 4 Ester	34704-124	43.4%	3.8	✓	✓	✓	N
Low Vol 6 Ester	34704-125	58.9%	5.6	✓	✓	✓	N
Saber	34704-803	38.7%	3.8	✓	✓	✓	N
Salvo Low Volatile Weed Killer	34704-609	54.2%	5.0	✓	✓	✓	N
Savage Dry Soluble	34704-606	78.9%	DS	✓	✓	✓	Y*
<b><u>Van Diest Supply (11773)</u></b>							
Cornbelt 4# Amine	11773-2	38.4%	3.8	✓	✓	✓	Y*
Cornbelt 4# LV Ester	11773-3	43.9%	3.8	✓	✓	✓	N
Cornbelt 6# LV Ester	11773-4	58.3%	5.6	✓	✓	✓	N
<b><u>Wilber-Ellis (2935)</u></b>							
Amine 4A Selective Herbicide	2935-512	39.3%	3.8	✓	✓	✓	Y*
LV 4	2935-511	44.3%	3.8	✓	✓	✓	N
LV 6	228-95-2935	87.3%	5.5	✓	✓	✓	N
LV 6 Low Volatile Herbicide	42750-20-2935	57.4%	5.5	✓	✓	✓	N

\* Denotes ground and air application.

## Summary SITE and USE RESTRICTIONS

	Grazing Restriction	Haying Restriction	Non-Agricultural Use	Right-of-Way	Aquatic Use	Aerial Application
Tordon (picloram)	1 qt/A or more Lactating dairy-14 d Non-lactating dairy & beef - none <sup>1/</sup> Slaughter interval-3 d	1 qt/A or more - 14 d	Yes	Yes	No	Yes
Milestone (aminopyralid)	None	14 d	Yes	Yes	No	Yes
ForeFront (aminopyralid+ 2,4-D)	None	14 d	Yes	No	No	Yes
2,4-D ester	Lactating dairy-7 d Slaughter interval-3 d	30 d	Yes	Yes	Some <sup>2/</sup>	Yes
2,4-D amine	Lactating dairy-7 d Slaughter interval-3 d	30 d	Yes	Yes	Some <sup>2/</sup>	Yes
Grazon P+D (picloram+ 2,4-D amine)	Lactating dairy-14 d Other livestock: no restrictions	30 d	No	No	No	Yes
Plateau (imazapic)	None	Do not harvest-7 d	Yes	Yes	No	Yes
Banvel, Clarity, Sterling (dicamba)	Lactating dairy: 1 pt/A-7 d 1 qt/A-21 d 2 qt/A-40 d Non-lactating dairy & beef-none Slaughter interval-30 d	Lactating dairy: 1 pt/A-37 d 1 qt/A-51 d 2 qt/A-70 d	Yes	Yes	No	Yes
Overdrive (diflufenzopyr+ dicamba)	None	None	Yes	Yes	No	CRP only
Stinger (clopyralid)	None	None	Yes	* <sup>2/</sup>	No	No
Transline (clopyralid)	None	None	Yes	Yes	No	Yes
Curtail (clopyralid+2,4-D)	Lactating dairy-14d Non-lactating dairy- none Slaughter interval-7 d <sup>3/</sup>	30 d	Yes	No	No	Yes
Ally or Cimarron (metsulfuron)	None	Use not listed	No	No	No	Yes
Ally, Cimarron+2,4-D (metsulfuron+2,4-D)	Lactating dairy-7 d Slaughter interval-30 d	Lactating dairy-37 d	No	No	No	Yes
Cimarron Max (metsulfuron+ dicamba+2,4-D)	Lactating dairy-7 d Slaughter interval-30 d	Lactating dairy-37 d	No	No	No	Yes
Escort XP (metsulfuron)	>1.66 oz-3 d	>1.66 oz-3 d	Yes	Yes	No	No
Telar (chlorsulfuron)	None for <1.3 oz	Use not listed	Yes	Yes	No	No
Arsenal (imazapyr)	Do not graze	Do not harvest	Yes	Yes	No	Yes
Habitat, Stalker (imazapyr)	Slaughter interval - 30 d	30 d	Yes	No	Yes	Helicopter Only

	Grazing Restriction	Haying Restriction	Non-Agricultural Use	Right-of-Way	Aquatic Use	Aerial Application
Krenite ( <i>fosamine</i> )	Do not graze-1 yr	Do not harvest-1 yr	Yes	Yes	No <sup>4/</sup>	Yes
Garlon 4L ( <i>triclopyr</i> )	Slaughter interval-3 d Do not graze lactating dairy until next growing season	Do not harvest-14 d	Yes	Yes	No	Yes
Garlon 3A ( <i>triclopyr</i> )			Yes	Yes	Labeled restricted areas	Yes
Krenite ( <i>fosamine</i> )	Do not graze-1 yr	Do not harvest-1 yr	Yes	Yes	No <sup>4/</sup>	Yes
Crossbow ( <i>triclopyr+2,4-D</i> )	Lactating dairy: <2 gal/A-14 d >2 gal/A-next year Other livestock: <2 gal/A-none 2-4 gal/A-14 d <sup>5/</sup> Slaughter interval-3 d	Lactating dairy: next year Other livestock: <2 gal/A-7 d >2 gal/A-14 d Removal before slaughter-3 d	Yes	Yes	No	Yes
Redeem ( <i>triclopyr+ clopyralid</i> )	Lactating dairy: 14 d Other livestock: No restriction	Lactating dairy: Do not harvest until next growing season. Other livestock: 7 d	Yes	No (fence-lines)	No	Yes
Paramount ( <i>quinclorac</i> )	Do not graze	Do not harvest for 20 days	Yes	Yes	No	No
Glyphosate Products	Slaughter interval-3 d Spot treatment <sup>6/</sup> -14 d Broadcast treatment-8 weeks	Spot treatment-14 d Broadcast treatment-8 wks	No	No	No	Yes
Roundup Pro	Do not graze	Do not harvest	Yes	Yes	No	Yes
Rodeo ( <i>glyphosate</i> )	----	----	Yes	Yes	Yes	Yes
Touchdown Pro ( <i>glyphosate</i> )	Do not graze	Do not harvest	Yes	Yes	No	Yes

<sup>1/</sup> Remove livestock to untreated grass pasture for 7 days before transferring livestock to broadleaf or broadleaf-mixed pasture areas. Otherwise, urine may contain enough product to cause injury to sensitive broadleaf plants.

<sup>2/</sup> Refer to specific label for aquatic use products.

<sup>3/</sup> Withdrawal not needed if 2 weeks or more time elapsed since application.

<sup>4/</sup> Apply only to water's edge.

<sup>5/</sup> If less than 25% of grazed area is treated, there is no grazing restrictions.

<sup>6/</sup> Do not treat more than 1/10 of any given acre at one time with spot or wiper applications. Remove livestock before application.

<sup>7/</sup> Fence rows, around farm buildings, and equipment pathways.



## **CHEMICAL, PHYSICAL and SAFETY CHARACTERISTICS of HERBICIDES**

Properties of the most commonly used herbicides to control noxious weeds are listed in the table below. Formulation and other local conditions will affect values for most properties. Solubility is affected by formulation, temperature and soil pH.

Half-life refers to the days required for the herbicide level in the soil to be reduced to half the original amount applied. Rainfall, temperature, and soil pH are important factors affecting half life.

Toxicity for oral ingestion and dermal exposure are expressed as the quantity required for a lethal dose for 50% of a population. The LD<sub>50</sub> value may be multiplied by .003 to determine ounces of active ingredient that would be lethal for half a population of 180 lb subjects.

Leaching (LE) rating refers to risk of herbicide movement through the soil profile into groundwater. Chemical properties of the herbicide, soil properties and rainfall are important factors affecting leaching potential. Herbicide movement in runoff solution (SL) is a rating for risk of movement in surface water. Rainfall amount and intensity, soil properties, surface characteristics, and herbicide rate are important variables affecting runoff solution.

Special safety equipment is based on label statements. All herbicides should be handled according to label safety guidelines using equipment and precautions that minimize risk of exposure.

Herbicide	Solubility (ppm)	Half Life (days)	Surface and Groundwater Risk		LD <sub>50</sub> (mg/kg)	
			Leaching (LE)	Runoff Sol. (SL)	Oral	Dermal
Tordon 22K (picloram)	200,000	90	High	High	8200	>2000
Milestone (aminopyralid)	2,480	35	Low	Low	>5000	>5000
2,4-D ester	8	7	Low	Low	375	800
2,4-D amine	796,000	10	Low	Low	375	800
Banvel (dicamba)	400,000	14	Low to Medium	Low	1707	>2000
Stinger (clopyralid)	300,000	30	Moderate Potential	Low	>5000	>2000
Roundup (glyphosate)	400,000	47	Low	Low	4300	>940
Arsenal (imazapyr)	11,000	90	Low	Low	>5000	>2148
Krenite (fosamine)	1,790,000	8	Low	Low	>5000	>5000
Garlon (triclopyr ester)	23	46	Medium	Medium	630	2140
Escort/Ally (metsulfuron)	9500	120	Moderate@high pH	Medium	>5000	<2000
Telar (chlorsulfuron)	7000	160	Moderate@high pH	Medium	5545	>3400
Plateau (imazapic)	2232	120	Low	Low	>5000	>5000

Data base: National Water Quality Technology Staff and other references.

## **BIOCONTROL of NOXIOUS WEEDS**

Biocontrol offers another tool for noxious weed control. The basic idea behind biocontrol is utilizing the weed pest's natural enemies as a means of weakening or killing the host plant. Insects used as biocontrol agents for noxious weeds include 1) seed attackers, 2) gall formers, 3) defoliators, 4) sap suckers, 5) stem dwellers, 6) crown and root burrowers, and 7) root feeders. Biocontrol of noxious weeds is not the best choice in all situations. If the noxious weed infestation can be controlled by a more efficient means, then that should be the option of choice. There are potential economic benefits as well as advantages where environmental situations or site restrictions limit or preclude other control options. Biocontrol approaches usually require at least 3 to 5 years for significant results. It is important to realize that a biological control program will not eradicate noxious weeds. A residual level of the weed population is to be expected even under the best conditions. Survival of the biocontrol agents is dependent on the density of the host noxious weeds. This is a natural cycle and should be expected so a resurgence of noxious weed populations may occur due to 1) seed reserves in the soil; 2) missed plants, and 3) lagging populations of the biocontrol agent. Ultimately, the greatest benefit will be realized if biocontrol is integrated with other control measures.

Development of biocontrol approaches in the state continues. Prior to the initial release by USDA-APHIS biocontrol agents are tested and studied to ensure they are host-specific for the weed targeted. USDA-APHIS and the South Dakota Department of Agriculture are involved with the initial introductions and development of insectary sites for future distributions. County weed and pest boards as well as state and federal agencies who manage lands in South Dakota have released several insects that have been evaluated and approved for release as biocontrol agents.

Approved biocontrol agents are available through several reputable insectary businesses located in neighboring states. Prices and availability vary on each individual bioagent from year to year.

**LEAFY SPURGE.** Several insects have been evaluated as a biological control agent on leafy spurge in South Dakota. The black flea beetle (*Aphthona lacertosa*) has shown the best results. This insect is quite adaptable to much of the state's diverse climate and environmental conditions. Flea beetles are sun loving insects that prefer dry sites; however, the black flea beetle will tolerate cooler, shaded, and wetter sites.

Currently in South Dakota there is a collection and release program for the leafy spurge flea beetles. This program is coordinated by the South Dakota Department of Agriculture. The county weed and pest board is the local contact point for landowners or land managers considering the use of flea beetles on their leafy spurge infestations. South Dakota landowners can collect the black flea beetle (*A. lacertosa*) or the blackdot (brown) flea beetle (*A. nigriscutis*) free of charge on state supervised collection days, which are usually held the end of June.

Several other approved bioagents were evaluated and they include root feeding leafy spurge flea beetle species (*Aphthona nigriscutis*, *A. czwalinae*, *A. flava* and *A. cyarissiae*), a stem boring larvae (*Oberea erythrocephala*), and a tip gall midge (*Spurgia esulae*).

**MUSK THISTLE.** *Rhinocyllus conicus*, a seedhead weevil, was initially released more than 20 years ago. It can be found statewide in musk thistle infestations. Seed production is substantially reduced. A rosette weevil, *Trichosirocalus horridus*, has also shown promise.

**CANADA THISTLE.** Introduction of bioagents is relatively new for Canada thistle in South Dakota. Choice of successful biocontrol agents has been limited because there are a number of desirable plant species related to Canada thistle. There have been releases of *Hadroplatus litura* (stem mining weevil), *Larinus planus* (seedhead weevil), *Cassida rubiginosa* (defoliating beetle), and *Urophora cardui* (stem and shoot gall fly). The stem mining weevil has shown the most potential for decreasing Canada thistle patches. The gall fly has some effect on seed production and seems to work best when released in combination with the stem mining weevil. Both of these species have good winter survival and reproductive success. The musk thistle seed weevil (*Rhinocyllus conicus*) has been found in some Canada thistle infestations; however, they have a limited effect on stand reduction.

**BIENNIAL KNAPWEED.** The primary bioagent released for the control of biennial knapweeds is *Larinus minutus*, the knapweed flowerhead weevil. Other insects studied and released include *Agapeta zoegana*, a root boring moth; *Cyphocleonus achates*, a root boring weevil; and *Sphenoptera jugoslavica*, a root boring beetle.

**DALMATIAN TOADFLAX.** One of the more recent releases for control of Dalmatian toadflax is a root galling weevil, *Gymnetron antirrhini*. Most of these releases have been in the Pennington and Lawrence county areas.

**ST. JOHNSWORT.** *Chrysolina quadrigemina* (foliage feeding beetle) has shown promise and has been introduced in some western South Dakota counties.

**PURPLE LOOSESTRIFE.** *Galerucella californiensis* (leaf feeding beetle) has been introduced in wetland sites invaded by this weed. This insect has produced dramatic results in other states. A root mining weevil, *Hylobius transversovittatus*, and two seed weevil species, *Nanophyes spp.*, have also been released. The *Galerucella* species has been used extensively in South Dakota since the development of an insectary and rearing site at the South Dakota Human Service Center Trustee Unit in Yankton, S.D.

**SALT CEDAR.** Several insect bioagents have been evaluated and approved in some southwestern and western states. The Chinese leaf beetle (*Diorhabda enlongata*), a defoliator, has shown promise. It is currently being evaluated in South Dakota but has not been approved for general release.



# Common Biocontrol Agents for Noxious Weeds in South Dakota

## LEAFY SPURGE BIOAGENTS



Fig. 1 *Aphthona nigricutis* adult (brown flea beetle)




Fig. 2 *Aphthona lacertosa* adult (black flea beetle)



Fig. 3 *Aphthona lacertosa* flea beetles on leafy spurge



Fig. 4 South Dakota flea beetle collection/beetle sorter

## CANADA THISTLE BIOAGENTS



Fig. 5 *Hadroplatus litura* adult (Canada thistle stem mining weevil)



Fig. 6 *Hadroplatus litura* larvae in Canada thistle stem



Fig. 7 Mined Canada thistle stem—Hand County



Fig. 8 *Larinus planus* adult (Canada thistle seedhead weevil)



Fig. 9 *Urophora cardui* (Canada thistle gall fly)



Fig. 10 Gall on Canada thistle stem—Beadle County



Fig. 11 *Cassida rubiginosa* adult (defoliating beetle)

## MUSK THISTLE BIOAGENTS



Fig. 12 *Rhinocyllus conicus* adult (musk thistle seed weevil)



Fig. 13 Left: Normal (musk thistle seedhead); Right: infested musk thistle seedhead



Fig. 14 *Trichosivocalus horridus* adult (musk thistle rosette weevil)

## SPOTTED KNAPWEED BIOAGENTS



Fig. 15 *Larinus minutus* adult (flowerhead weevil)




Fig. 16 Knapweed seedhead with emergence hole—Pennington County